

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
7 September 2001 (07.09.2001)

PCT

(10) International Publication Number
WO 01/64655 A1(51) International Patent Classification⁷: C07D 239/48,
403/12

(21) International Application Number: PCT/GB01/00824

(22) International Filing Date: 26 February 2001 (26.02.2001)

(25) Filing Language: English

(26) Publication Language: English



(30) Priority Data:
0004890.0 1 March 2000 (01.03.2000) GB(71) Applicant (for all designated States except MG, US): AS-
TRAZENECA AB [SE/SE]; S-151 85 Sodertalje (SE).(71) Applicant (for MG only): ASTRAZENECA UK LIM-
ITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN
(GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PEASE, Elizabeth,
Janet [GB/GB]; Alderley Park, Macclesfield, Cheshire
SK10 4TG (GB). BREAUULT, Gloria, Anne [US/GB];Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).
WILLIAMS, Emma, Jane [GB/GB]; Alderley Park,
Macclesfield, Cheshire SK10 4TG (GB). BRADBURY,
Robert, Hugh [GB/GB]; Alderley Park, Macclesfield,
Cheshire SK10 4TG (GB). MORRIS, Jeffrey, James
[GB/GB]; Alderley Park, Macclesfield, Cheshire SK10
4TG (GB).(74) Agent: BRYANT, Tracey; AstraZeneca, Global Intellec-
tual Property, Patents, Alderley Park, Mereside, P.O. Box
272, Macclesfield, Cheshire SK10 4GR (GB).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: 2, 4-DI(HETERO-)ARYLAMINO (-OXY)-5-SUBSTITUTED PYRIMIDINES AS ANTINEOPLASTIC AGENTS



(57) Abstract: Pyrimidine derivatives of the formula (I), wherein: Q₁ and Q₂ are independently selected from aryl or carbon linked heteroaryl optionally substituted as defined within; and one or both Q₁ and Q₂ are substituted on a ring carbon by one substituent of the formula (Ia) or (Ia'), wherein: Y, Z, n, m, Q₃, G, R¹, are as defined within; and pharmaceutically acceptable salts and in vivo hydrolysable esters thereof are described. Processes for their manufacture, pharmaceutical compositions and their use as cyclin-dependent serine/threonine kinase (CDK) and focal adhesion kinase (FAK) inhibitors are also described.

WO 01/64655 A1



Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

2,4-DI (HETERO-) ARYLAMINO (-OXY) -5-SUBSTITUTED PYRIMIDINES AS ANTINEOPLASTIC AGENTS

The invention relates to pyrimidine derivatives, or pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof, which possess cell-cycle inhibitory activity and are accordingly useful for their anti-cancer (such as anti-cell-proliferative, anti-cell migration and/or apoptotic) activity and are therefore useful in methods of treatment of a warm-blooded animal, such as man. The invention also relates to processes for the manufacture of said pyrimidine derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments or use in the production of an anti-cancer (anti-cell-proliferation/migration and/or apoptotic) effect in a warm-blooded animal such as man.

A family of intracellular proteins called cyclins play a central role in the cell cycle. The synthesis and degradation of cyclins is tightly controlled such that their level of expression fluctuates during the cell cycle. Cyclins bind to cyclin-dependent serine/threonine kinases (CDKs) and this association is essential for CDK (such as CDK1, CDK2, CDK4 and/or CDK6) activity within the cell. Although the precise details of how each of these factors combine to regulate CDK activity is poorly understood, the balance between the two dictates whether or not the cell will progress through the cell cycle.

The recent convergence of oncogene and tumour suppresser gene research has identified regulation of entry into the cell cycle as a key control point of mitogenesis in tumours. Moreover, CDKs appear to be downstream of a number of oncogene signalling pathways. Disregulation of CDK activity by upregulation of cyclins and/or deletion of endogenous inhibitors appears to be an important axis between mitogenic signalling pathways and proliferation of tumour cells.

Accordingly it has been recognised that an inhibitor of cell cycle kinases, particularly inhibitors of CDK2, CDK4 and/or CDK6 (which operate at the S-phase, G1-S and G1-S phase respectively) should be of value as a selective inhibitor of cell proliferation, such as growth of mammalian cancer cells.


Furthermore, it is believed that inhibition of focal adhesion kinase (FAK), which is involved in signal transduction pathways, induces apoptosis (cell-death) and/or inhibits cell migration and an inhibitor of FAK may therefore have value as an anti-cancer agent.

The present invention is based on the discovery that certain 2,4-pyrimidine compounds surprisingly inhibit the effects of cell cycle kinases showing selectivity for CDK2, CDK4 and CDK6, and also inhibit FAK and thus possess anti-cancer (anti-cell-migration/proliferation and/or apoptotic) properties. Such properties are expected to be of value in the treatment of

5 disease states associated with aberrant cell cycles and cell proliferation such as cancers (solid tumours and leukemias), fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic inflammation, bone diseases and ocular diseases with retinal vessel proliferation.

10 According to the invention there is provided a pyrimidine derivative of the formula


(I):




(I)

wherein:

15 Q_1 and Q_2 are independently selected from aryl or carbon linked heteroaryl; and one of Q_1 and Q_2 or both of Q_1 and Q_2 is substituted on a ring carbon by one substituent of the formula (Ia) or (Ia'):



(Ia)



(Ia')

20 wherein:

Y is -NHC(O)- or -C(O)NH-;

Z is R^aO- , R^bR^cN- , R^dS- , $R^eR^fNNR^g-$, C_{3-8} cycloalkyl, phenyl or a heterocyclic group;

wherein said phenyl, C_{3-8} cycloalkyl or heterocyclic group are optionally substituted on a ring carbon by one or more groups selected from R^h ; and wherein if said heterocyclic group

contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^i ;

R^a , R^b , R^c , R^d , R^e , R^f and R^g are independently selected from hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl and C_{3-8} cycloalkyl; wherein said C_{1-4} alkyl, C_{2-4} alkenyl and C_{3-8} cycloalkyl are
5 optionally substituted by one or more groups selected from R^j ;

n is 0 or 1;

m is 1, 2 or 3;

Q_3 is a nitrogen linked heterocycle; wherein said heterocycle is optionally substituted on a ring carbon by one or more groups selected from R^k ; and wherein if said heterocyclic
10 group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^m ;

G is -O- or -NR²-;

R^2 is selected from hydrogen, C_{1-6} alkyl, C_{3-6} alkenyl and C_{3-6} alkynyl; wherein said C_{1-6} alkyl, C_{3-6} alkenyl and C_{3-6} alkynyl are optionally substituted by one or more groups
15 selected from R^n ;

R^1 is selected from hydrogen, halo, hydroxy, nitro, amino, N -(C_{1-3} alkyl)amino, N,N -di-(C_{1-3} alkyl)amino, cyano, trifluoromethyl, trichloromethyl, C_{1-3} alkyl [optionally substituted by 1 or 2 substituents independently selected from halo, cyano, amino, N -(C_{1-3} alkyl)amino, N,N -di-(C_{1-3} alkyl)amino, hydroxy and trifluoromethyl], C_{3-5} alkenyl
20 [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent], C_{3-5} alkynyl, C_{1-3} alkoxy, mercapto, C_{1-3} alkylsulphanyl, carboxy and C_{1-3} alkoxycarbonyl;

Q_1 is optionally substituted on a ring carbon by one to four substituents independently selected from halo, mercapto, nitro, formyl, formamido, carboxy, cyano, amino, ureido, carbamoyl, sulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl [wherein said C_{1-4} alkyl, C_{2-4} alkenyl
25 and C_{2-4} alkynyl are optionally substituted by one or more groups selected from R^o], C_{1-4} alkanoyl, C_{1-4} alkoxycarbonyl, heterocyclic group, C_{1-4} alkylS(O) _{a} wherein a is 0 to 2 [optionally substituted by hydroxy], N -(C_{1-4} alkyl)ureido, N,N -di-(C_{1-4} alkyl)ureido, N' -(C_{1-4} alkyl)- N -(C_{1-4} alkyl)ureido, N',N' -di-(C_{1-4} alkyl)- N -(C_{1-4} alkyl)ureido, N - C_{1-4} alkylamino, N,N -di-(C_{1-4} alkyl)amino, N -(C_{1-4} alkyl)sulphamoyl, N,N -di-(C_{1-4} alkyl)sulphamoyl,
30 N - C_{1-4} alkylcarbamoyl, N,N -di-(C_{1-4} alkyl)carbamoyl and C_{1-4} alkanoylamino;

and also independently, or in addition to, the above substituents, Q_1 may be optionally substituted by one to two substituents independently selected from aryl, C_{3-8} cycloalkyl and a

heterocyclic group; wherein said aryl, C₃₋₈cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^p; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^q;

- 5 and also independently, or in addition to, the above substituents, Q₁ may be optionally substituted by one C₁₋₄alkoxy or by one hydroxy substituent;

- Q₂ is optionally substituted on a ring carbon by one to four substituents independently selected from halo, hydroxy, mercapto, nitro, formyl, formamido, carboxy, cyano, amino, ureido, carbamoyl, sulphamoyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy [wherein said
- 10 C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl and C₁₋₄alkoxy are optionally substituted by one or more groups selected from R^j], C₁₋₄alkanoyl, C₁₋₄alkoxycarbonyl, heterocyclic group, C₁₋₄alkylS(O)_a wherein a is 0 to 2 [optionally substituted by hydroxy], Nⁱ-(C₁₋₄alkyl)ureido, Nⁱ,Nⁱ-di-(C₁₋₄alkyl)ureido, Nⁱ-(C₁₋₄alkyl)-N-(C₁₋₄alkyl)ureido, Nⁱ,Nⁱ-di-(C₁₋₄alkyl)-N-(C₁₋₄alkyl)ureido, N-C₁₋₄alkylamino, N,N-di-(C₁₋₄alkyl)amino,
- 15 N-(C₁₋₄alkyl)sulphamoyl, N,N-di-(C₁₋₄alkyl)sulphamoyl, N-C₁₋₄alkylcarbamoyl, N,N-di-(C₁₋₄alkyl)carbamoyl, C₂₋₄alkenyloxy, C₂₋₄alkynyloxy, C₁₋₄alkanoylamino and a group of formula (Ia) or (Ia') as depicted above;

- and also independently, or in addition to, the above substituents, Q₂ may be optionally substituted by one to two substituents independently selected from aryl, C₃₋₈cycloalkyl or a
- 20 heterocyclic group; wherein said aryl, C₃₋₈cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^s; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^t;

- Rⁱ, R^o, R^o and R^r are independently selected from hydroxy, halo, amino, cyano,
- 25 formyl, formamido, carboxy, nitro, mercapto, carbamoyl, sulphamoyl, N-C₁₋₄alkylamino, N,N-di-(C₁₋₄alkyl)amino, C₁₋₄alkanoyl, C₁₋₄alkanoyloxy, C₁₋₄alkoxy, C₁₋₄alkoxycarbonyl, N-C₁₋₄alkylcarbamoyl, N,N-di-(C₁₋₄alkyl)carbamoyl, C₁₋₄alkanoylamino, C₁₋₄alkylS(O)_a wherein a is 0 to 2, C₁₋₄alkylsulphonylamino, N-(C₁₋₄alkyl)sulphamoyl, N-(C₁₋₄alkyl)₂sulphamoyl, N-(C₁₋₄alkyl)carbamoyl, N-(C₁₋₄alkyl)₂carbamoyl, phenyl,
- 30 phenylthio, phenoxy, C₃₋₈cycloalkyl and a heterocyclic group; wherein said phenyl, phenylthio, phenoxy, C₃₋₈cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^u; and wherein if said heterocyclic group

contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^v;

R^b, R^k, R^p, R^t and R^u are independently selected from hydroxy, halo, amino, cyano, formyl, formamido, carboxy, nitro, mercapto, carbamoyl, sulphamoyl, C₁₋₄alkyl [optionally substituted by one or more groups selected from halo, cyano, amino, *N*-C₁₋₄alkylamino, *N,N*-di-(C₁₋₄alkyl)amino or hydroxy], C₂₋₄alkenyl [optionally substituted by one or more groups selected from halo], C₂₋₄alkynyl, *N*-C₁₋₄alkylamino, *N,N*-di-(C₁₋₄alkyl)amino, C₁₋₄alkanoyl, C₁₋₄alkanoyloxy, C₁₋₄alkoxy [optionally substituted by one or more groups selected from halo], C₁₋₄alkoxycarbonyl, *N*-C₁₋₄alkylcarbamoyl, *N,N*-di-(C₁₋₄alkyl)carbamoyl, C₁₋₄alkanoylamino, C₁₋₄alkylS(O)_a, wherein a is 0 to 2, C₁₋₄alkylsulphonylamino, *N*-(C₁₋₄alkyl)sulphamoyl, *N*-(C₁₋₄alkyl)₂sulphamoyl, phenyl, C₃₋₈cycloalkyl and a heterocyclic group; and

Rⁱ, R^q, R^r and R^v are independently selected from C₁₋₄alkyl, C₁₋₄alkanoyl, C₁₋₄alkylsulphonyl, C₁₋₄alkoxycarbonyl, carbamoyl, *N*-(C₁₋₄alkyl)carbamoyl, *N,N*-(C₁₋₄alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

"Aryl" is a fully or partially unsaturated, mono or bicyclic carbon ring that contains 4-12 atoms. Preferably "aryl" is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. More preferably "aryl" is phenyl, naphthyl, tetralinyl or indanyl. Particularly "aryl" is phenyl, naphthyl or indanyl. More particularly "aryl" is phenyl.

A "carbon linked heteroaryl" is a fully unsaturated, 5- or 6- membered monocyclic ring or 9- or 10- membered bicyclic ring of which at least one atom is chosen from nitrogen, sulphur or oxygen. This ring is linked via a carbon atom to the -NH- (for Q₁) or G (for Q₂). Preferably "carbon linked heteroaryl" is furanyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyrazolyl, furazanyl, triazolyl, thiadiazolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazinyl, indolyl, quinolyl or benzimidazolyl. More preferably "carbon linked heteroaryl" is pyridyl, thiazolyl or pyrazolyl. Particularly "carbon linked heteroaryl" is pyridyl.

A "heterocyclic group" is a saturated, partially saturated or unsaturated, mono or bicyclic ring containing 4-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a -CH₂- group can optionally be replaced by a -C(O)-, and a ring sulphur atom may

be optionally oxidised to form S-oxide(s). Preferably a "heterocyclic group" is pyrrolidinyl, morpholino, piperidyl, pyridyl, pyranyl, pyrrolyl, isothiazolyl, indolyl, quinolyl, thienyl, furyl, 1,3-benzodioxolyl, thiadiazolyl, piperazinyl, thiazolidinyl, pyrrolidinyl, thiomorpholino, pyrazolyl, pyrrolinyl, homopiperazinyl, tetrahydropyranyl, imidazolyl, pyrimidyl, pyrazinyl, 5 pyridazinyl, isoxazolyl, 4-pyridone, 1-isoquinolone, 2-pyrrolidone, 4-thiazolidone, imidazo[1,2-a]pyridine or 3-aza-8-oxabicyclo[3,2,1]hexane. More preferably a "heterocyclic group" is pyrrolidinyl, morpholino, piperidyl, indolyl, thienyl, furyl, piperazinyl, thiomorpholino, pyrazolyl, imidazolyl, 2-pyrrolidone, imidazo[1,2-a]pyridine or 3-aza-8-oxabicyclo[3,2,1]hexane.

- 10 A "nitrogen linked heterocycle" is a saturated, partially saturated or fully unsaturated, mono or bicyclic ring containing 4-12 atoms, one atom of which is a nitrogen atom (attached to form an amide as shown) and the other atoms are either all carbon atoms or they are carbon atoms and 1-3 heteroatoms chosen from nitrogen, sulphur or oxygen, wherein a -CH₂- group can optionally be replaced by a -C(O)- and a ring sulphur atom may be optionally oxidised to 15 form the S-oxides. It will be appreciated that in forming this nitrogen link, the nitrogen atom is not quaternised, i.e. a neutral compound is formed. Preferably "nitrogen linked heterocycle" is pyrrol-1-yl, pyrrolin-1-yl, pyrrolidin-1-yl, imidazol-1-yl, imidazolin-1-yl, imidazolidin-1-yl, pyrazol-1-yl, pyrazolin-1-yl, pyrazolidin-1-yl, triazol-1-yl, piperidin-1-yl, piperazin-1-yl, morpholino, thiomorpholino, indol-1-yl, indolidin-1-yl or benzimidazol-1-yl. 20 More preferably "nitrogen linked heterocycle" is piperidin-1-yl.

In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. An analogous convention applies to other generic terms. "Halo" is fluoro, chloro, bromo and iodo.

- 25 Examples of C₂₋₄alkenyl are vinyl and allyl; examples of C₂₋₆alkenyl are C₃₋₅alkenyl, vinyl and allyl; an example of C₃₋₆alkenyl is allyl; an examples of C₃₋₆alkynyl are C₃₋₅alkynyl and propyn-2-yl; examples of C₂₋₄alkynyl are ethynyl and propyn-2-yl; examples of C₂₋₆alkynyl are ethynyl and propyn-2-yl; examples of C₁₋₄alkanoyl are acetyl and propionyl; examples of C₁₋₄alkoxycarbonyl are C₁₋₃alkoxycarbonyl, methoxycarbonyl, ethoxycarbonyl, 30 propoxycarbonyl and *tert*-butoxycarbonyl; examples of C₁₋₄alkylene are methylene, ethylene and propylene; examples of C₁₋₄alkyl are C₁₋₃alkyl, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl and *tert*-butyl; examples of C₁₋₆alkyl are methyl, ethyl, propyl, isopropyl,

- butyl, isobutyl, *sec*-butyl, *tert*-butyl and 3-methylbutyl; examples of C_{1-4} alkoxy are C_{1-3} alkoxy, methoxy, ethoxy, propoxy, isopropoxy and butoxy; an example of C_{2-4} alkenyloxy is allyloxy; an example of C_{2-4} alkynyloxy is propynyloxy; examples of C_{1-4} alkylS(O)_a, wherein a is 0 to 2 are C_{1-3} alkylsulphanyl, methylthio, ethylthio, propylthio, methylsulphinyl, ethylsulphinyl, propylsulphinyl, mesyl, ethylsulphonyl and propylsulphonyl; examples of N - C_{1-4} alkylcarbamoyl are N -methylcarbamoyl, N -ethylcarbamoyl and N -propylcarbamoyl; examples of N,N -di-(C_{1-4} alkyl)-carbamoyl are N,N -dimethylcarbamoyl, N -ethyl- N -methylcarbamoyl and N,N -diethylcarbamoyl; examples of N - C_{1-4} alkylamino are N -(C_{1-3} alkyl)amino, methylamino, ethylamino and propylamino; examples of N,N -di-(C_{1-4} alkyl)amino are N,N -di-(C_{1-3} alkyl)amino, dimethylamino, N -ethyl- N -methylamino, diethylamino, N -methyl- N -propylamino and dipropylamino; examples of C_{1-4} alkanoylamino are acetamido, propionamido and butyramido; examples of C_{3-8} cycloalkyl are cyclopropyl, cyclopentyl and cyclohexyl; examples of C_{1-4} alkanoyl are acetyl and propionyl; examples of C_{1-4} alkanoyloxy are acetyloxy and propionyloxy; examples of N' -(C_{1-4} alkyl)ureido are N' -methylureido and N' -ethylureido; examples of N',N' -di-(C_{1-4} alkyl)ureido are N',N' -dimethylureido, N',N' -diisopropylureido and N' -methyl- N' -propylureido; examples of N' -(C_{1-4} alkyl)- N -(C_{1-4} alkyl)ureido are N' -methyl- N -ethylureido and N' -methyl- N -methylureido; examples of N',N' -di-(C_{1-4} alkyl)- N -(C_{1-4} alkyl)ureido are N',N' -dimethyl- N -ethylureido and N' -methyl- N' -propyl- N -butylureido; examples of N -(C_{1-4} alkyl)sulphamoyl are N -methylsulphamoyl and N -isopropylsulphamoyl; examples of N,N -di-(C_{1-4} alkyl)sulphamoyl are N -methyl- N -ethylsulphamoyl and N,N -dipropylsulphamoyl; and examples of C_{1-4} alkylsulphonylamino are mesylamino, ethylsulphonylamino and propylsulphonylamino.
- 25 A suitable pharmaceutically acceptable salt of a pyrimidine derivative of the invention is, for example, an acid-addition salt of a pyrimidine derivative of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In addition a suitable pharmaceutically acceptable salt of a pyrimidine derivative of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically
- 30

acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

The compounds of the formula (I) may be administered in the form of a pro-drug which is broken down in the human or animal body to give a compound of the formula (I).

5 Examples of pro-drugs include *in vivo* hydrolysable esters of a compound of the formula (I).

An *in vivo* hydrolysable ester of a compound of the formula (I) containing carboxy or hydroxy group is, for example, a pharmaceutically acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically acceptable esters for carboxy include C₁₋₆alkoxymethyl esters for example methoxymethyl,

10 C₁₋₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

15 An *in vivo* hydrolysable ester of a compound of the formula (I) containing a hydroxy group includes inorganic esters such as phosphate esters and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxy-methoxy. A selection of *in vivo* hydrolysable ester forming groups
20 for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and *N*-(dialkylaminoethyl)-*N*-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl. Examples of substituents on benzoyl include morpholino and piperazino linked from a ring nitrogen atom via a methylene group to the 3- or 4- position of the benzoyl ring.

25 Some compounds of the formula (I) may have chiral centres and/or geometric isomeric centres (E- and Z- isomers), and it is to be understood that the invention encompasses all such optical, diastereo-isomers and geometric isomers that possess CDK and/or FAK inhibitory activity.

The invention relates to any and all tautomeric forms of the compounds of the formula
30 (I) that possess CDK and/or FAK inhibitory activity.

It is also to be understood that certain compounds of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be

understood that the invention encompasses all such solvated forms which possess CDK and/or FAK inhibitory activity.

Particular preferred compounds of the invention comprise a pyrimidine derivative of the formula (I), or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof,
5 wherein R¹, Q₁, Q₂, and G have any of the meanings defined hereinbefore, or any of the following values. Such values may be used where appropriate with any of the definitions, claims or embodiments defined hereinbefore or hereinafter.

Preferably Q₁ and Q₂ are independently selected from phenyl, pyridyl, thiazolyl and pyrazolyl.

10 Preferably Q₁ is phenyl.

Preferably Q₂ is phenyl, pyridyl, thiazolyl or pyrazolyl.

More preferably Q₂ is phenyl or pyridyl.

Preferably Q₁ is phenyl and Q₂ is selected from phenyl, pyridyl, thiazolyl and pyrazolyl.

15 More preferably Q₁ is phenyl and Q₂ is selected from phenyl and pyridyl.

Preferably in the substituent (Ia) or (Ia'):

Y is -NHC(O)- or -C(O)NH-;

Z is R^aO-, R^bR^cN-, R^dS-, phenyl or a heterocyclic group; wherein said phenyl or heterocyclic group are optionally substituted on a ring carbon by one or more groups selected
20 from R^b; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from Rⁱ;

R^a, R^b, R^c and R^d are C₁₋₄alkyl;

n is 0 or 1;

m is 1, 2 or 3; and

25 Q₃ is a nitrogen linked heterocycle; wherein said heterocycle is optionally substituted on a ring carbon by one or more groups selected from R^k.

More preferably in the substituent (Ia) or (Ia'):

Y is -NHC(O)- or -C(O)NH-;

Z is R^aO-, R^bR^cN-, R^dS-, phenyl, pyrrolidinyl, morpholino, piperidyl, indolyl, thienyl,
30 furyl [optionally substituted by one or more methyl], piperazinyl [optionally substituted on a ring nitrogen by methyl], thiomorpholino, pyrazolyl [optionally substituted by one or more methyl], imidazolyl [optionally substituted by one or more groups selected from groups

selected from methyl and nitro], 2-pyrrolidone, imidazo[1,2-a]pyridine or 3-aza-8-oxabicyclo[3,2,1]hexane;

R^a , R^b , R^c and R^d are C_{1-4} alkyl;

n is 0 or 1;

5 m is 1, 2 or 3; and

Q_3 is piperidin-1-yl optionally substituted by pyrrolidin-1-yl.

Particularly the substituent **(Ia)** or **(Ia')** is

- N*-[2-(3-aza-8-oxabicyclo[3,2,1]hex-3-yl)ethyl]carbamoyl,
N-(2-di-*n*-butylaminoethyl)carbamoyl, *N*-(2-diethylaminoethyl)carbamoyl,
 10 *N*-(2-diisopropylaminoethyl)carbamoyl, *N*-[2-(3,5-dimethylpyrazol-1-yl)ethyl]carbamoyl,
N-(2-indol-3-ylethyl)carbamoyl, *N*-(2-isopropylaminoethyl)carbamoyl,
N-[2-(2-methyl-5-nitroimidazol-1-yl)ethyl]carbamoyl, *N*-(2-methylthioethyl)carbamoyl,
N-(2-morpholinoethyl)carbamoyl, *N*-(2-piperidin-1-ylethyl)carbamoyl,
N-(2-pyrrolidin-1-ylethyl)carbamoyl, *N*-(2-thien-2-ylethyl)carbamoyl,
 15 *N*-(2-thiomorpholinoethyl)carbamoyl, *N*-(3-*n*-butoxypropyl)carbamoyl,
N-(3-di-*n*-butylaminopropyl)carbamoyl, *N*-(3-imidazol-1-ylpropyl)carbamoyl,
N-[3-(4-methylpiperazin-1-yl)propyl]carbamoyl, *N*-(3-methylthiopropyl)carbamoyl,
N-(3-morpholinopropyl)carbamoyl, *N*-[3-(2-oxpyrrolidin-1-yl)propyl]carbamoyl,
N-(3-pyrrolidin-1-ylpropyl)carbamoyl, *N*-(2-di-isopropylaminoethyl)carbamoylmethyl,
 20 *N*-(2-morpholinoethyl)carbamoylmethyl, *N*-(4-dimethylaminobenzylamine)carbamoyl,
N-(imidazo[1,2-a]pyrid-2-ylmethyl)carbamoyl, *N*-(5-methylfur-2-ylmethyl)carbamoyl,
 4-(pyrrolidin-1-yl)piperidin-1-ylcarbonyl, 3-(dimethylamino)propanamide or
 3-(isopropylamino)propanamide.

More particularly the substituent **(Ia)** or **(Ia')** is *N*-(2-isopropylaminoethyl)carbamoyl,

- 25 *N*-(2-pyrrolidin-1-ylethyl)carbamoyl or *N*-(3-imidazol-1-ylpropyl)carbamoyl.

Preferably the substituent of formula **(Ia)** or **(Ia')** is in ring Q_1 .

Preferably when Q_1 is phenyl the substituent of formula **(Ia)** or **(Ia')** is in either the para- or meta- position relative to the -NH-.

- More preferably when Q_1 is phenyl the substituent of formula **(Ia)** or **(Ia')** is in the
 30 para- position relative to the -NH-.

In one aspect of the invention preferably G is -O-.

In a further aspect of the invention preferably G is -NR²-.

In one aspect of the invention when G is -NR²-, preferably R² is hydrogen.

In another aspect of the invention when G is -NR²-, preferably R² is not hydrogen.

Preferably R¹ is fluoro, chloro, bromo, methyl or cyano.

More preferably R¹ bromo.

5 Preferably Q₁ is unsubstituted except by the substituent of formula (Ia) or (Ia').

Preferably Q₂ is unsubstituted or substituted by one or two groups selected from fluoro, bromo, methyl, methoxy, methylthio or hydroxymethyl.

More preferably Q₂ is unsubstituted or substituted by one or two groups selected from fluoro, methyl or hydroxymethyl.

10 Preferably Q₂ is phenyl, 2-hydroxymethylphenyl, 3-fluorophenyl, 3-methylphenyl, 4-fluorophenyl, 4-bromophenyl, 4-methoxyphenyl, 4-hydroxymethylphenyl, 4-methylthiophenyl, 3,4-difluorophenyl, pyrid-2-yl, 6-methylpyrid-2-yl, 4-methylthiazol-2-yl or 5-methylpyrazol-2-yl.

More preferably Q₂ is pyrid-2-yl, 6-methylpyrid-2-yl, 3-fluorophenyl or
15 2-hydroxymethylphenyl.

Therefore, in a preferred aspect of the invention there is provided a pyrimidine derivative of the formula (I) as depicted above, wherein:

Q₁ and Q₂ are independently selected from phenyl, pyridyl, thiazolyl and pyrazolyl; Q₂ is unsubstituted or substituted by one or two groups selected from fluoro, bromo, methyl, methoxy, methylthio or hydroxymethyl; and Q₁ is substituted on a ring carbon by a
20 substituent of formula (Ia) or (Ia') as depicted above wherein:

Y is -NHC(O)- or -C(O)NH-;

Z is R^aO-, R^bR^cN-, R^dS-, phenyl or a heterocyclic group; wherein said phenyl or heterocyclic group are optionally substituted on a ring carbon by one or more groups selected
25 from R^h; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from Rⁱ;

R^a, R^b, R^c and R^d are C₁₋₄alkyl;

n is 0 or 1;

m is 1, 2 or 3; and

30 Q₃ is a nitrogen linked heterocycle; wherein said heterocycle is optionally substituted on a ring carbon by one or more groups selected from R^k.

G is -O- or -NH-; and

R¹ is fluoro, chloro, bromo, methyl or cyano;
or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

Therefore, in a more preferred aspect of the invention there is provided a pyrimidine derivative of the formula (I) as depicted above, wherein:

- 5 Q₁ is phenyl and Q₂ is pyrid-2-yl, 6-methylpyrid-2-yl, 3-fluorophenyl or 2-hydroxymethylphenyl; and Q₁ is substituted para to the -NH- linker by a group selected from *N*-(2-isopropylaminoethyl)carbamoyl, *N*-(2-pyrrolidin-1-ylethyl)carbamoyl or *N*-(3-imidazol-1-ylpropyl)carbamoyl;

G is -NH-; and

- 10 R¹ is bromo;
or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

In one aspect of the invention preferred compounds of the invention are those of Examples 41, 80, 82, 84, 85, 92, 94, 96, 114 or 115 or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

- 15 In a further aspect of the invention preferred compounds of the invention include any one of the Examples or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

Preferred aspects of the invention are those which relate to the compound or a pharmaceutically acceptable salt thereof.

- A pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or an
20 *in vivo* hydrolysable ester thereof, may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Such processes, when used to prepare a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, are provided as a further feature of the invention and are illustrated by the following representative examples in which, unless otherwise stated R¹, Q₁, Q₂ and G
25 have any of the meanings defined hereinbefore for a pyrimidine derivative of the formula (I) and unless another substituent is drawn on ring Q₁ or Q₂ the ring may bear any of the substituents described hereinbefore (optionally protected as necessary). Where a substituent is drawn on ring Q₁, this includes (unless stated otherwise) the possibilities of the substituent being on ring Q₂ in addition to, or instead of the substituent being on ring Q₁. Necessary
30 starting materials may be obtained by standard procedures of organic chemistry (see for example, Advanced Organic Chemistry (Wiley-Interscience), Jerry March - also useful for general guidance on reaction conditions and reagents). The preparation of such starting


materials is described within the accompanying non-limiting processes and Examples.

Alternatively necessary starting materials are obtainable by analogous procedures to those illustrated which are within the ordinary skill of an organic chemist.

Thus, as a further feature of the invention there are provided the following processes


5 which comprises of:-

a) for compounds of formula (I) where G is $-NR^2-$; reacting a pyrimidine of formula (II):



(II)


wherein L is a displaceable group as defined below, with a compound of formula (III):



(III)


where G is $-NR^2-$;

b) reaction of a pyrimidine of formula (IV):



(IV)


wherein L is a displaceable group as defined below, with a compound of formula (V):



(V)


c) for compounds of formula (I) wherein the sidechain is of formula (Ia) and Y is $-C(O)NH-$;

20 by reaction of an acid of formula (VI):



(VI)


or an activated derivative thereof; with an amine of formula (VII):



(VII)


5

d) for compounds of formula (I) wherein the sidechain is of formula (Ia) and Y is -NHC(O)- by reaction of an amine of formula (VIII):



(VIII)

10 with an acid of formula (IX):




(IX)

or an activated derivative thereof;

e) for compounds of formula (I) wherein the sidechain is of formula (Ia'); by reaction of an

15 acid of formula (VI) (or an activated derivative thereof) with an amine of formula (X):



(X)

and thereafter if necessary:

i) converting a compound of the formula (I) into another compound of the formula (I);

20 ii) removing any protecting groups;

iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

L is a displaceable group, suitable values for L are for example, a halo, sulphonyloxy or sulphur group, for example a chloro, bromo, methanesulphonyloxy, toluene-4-sulphonyloxy, mesyl, methylthio and methylsulphinyl.


25 Specific reaction conditions for the above reactions are as follows:-

Process a)

Pyrimidines of formula (II) and compounds of formula (III) may be reacted together:

- i) optionally in the presence of a suitable acid, for example an inorganic acid such as hydrochloric acid or sulphuric acid, or an organic acid such as acetic acid or formic acid. The reaction is preferably carried out in a suitable inert solvent or diluent, for example dichloromethane (DCM), acetonitrile, butanol, tetramethylene sulphone, tetrahydrofuran, 1,2-dimethoxyethane, *N,N*-dimethylformamide, *N,N*-dimethylacetamide or *N*-methylpyrrolidin-2-one, and at a temperature in the range, for example, 0° to 150°C, conveniently at or near reflux temperature; or
- ii) under standard Buchwald conditions (for example see *J. Am. Chem. Soc.*, **118**, 7215; *J. Am. Chem. Soc.*, **119**, 8451; *J. Org. Chem.*, **62**, 1568 and 6066) for example in the presence of palladium acetate, in a suitable solvent for example an aromatic solvent such as toluene, benzene or xylene, with a suitable base for example an inorganic base such as caesium carbonate or an organic base such as potassium-*t*-butoxide, in the presence of a suitable ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl and at a temperature in the range of 25 to 80°C.

Pyrimidines of the formula (II) may be prepared according to the following scheme:




wherein R^a is an optionally substituted alkyl or aryl group and L is a displaceable group as defined above. Preferably R^a is methyl, ethyl or *p*-tolyl.

Compounds of formula (III) are commercially available or are prepared by processes known in the art.

Process b)

- Pyrimidines of formula (IV) and anilines of formula (V) may be reacted together, i) in the presence of a suitable solvent for example a ketone such as acetone or an alcohol such as ethanol or butanol or an aromatic hydrocarbon such as toluene or *N*-methyl pyrrolidine, optionally in the presence of a suitable acid such as those defined above (or a suitable Lewis acid) and at a temperature in the range of 0°C to reflux, preferably reflux; or ii) under standard Buchwald conditions as described above.

- 10 Pyrimidines of formula (IV) are prepared according to the following scheme:



wherein L is a displaceable group as defined above.

The anilines of formula (V) are commercially available or are prepared by processes known in the art.


- 15 Pyrimidines of the formula (IVA) are commercially available or may be prepared by, for example, reacting a compound of formula (IVA) in which L is -OH (i.e. a uracil), with POCl₃ to give a compound of formula (IVA) in which L is -Cl.

Process c)

- Acids of formula (VI) and amines of formula (VII) may be coupled together in the presence of a suitable coupling reagent. Standard peptide coupling reagents known in the art can be employed as suitable coupling reagents, or for example carbonyldiimidazole and dicyclohexyl-carbodiimide, optionally in the presence of a catalyst such as dimethylaminopyridine or 4-pyrrolidinopyridine, optionally in the presence of a base for example triethylamine, pyridine, or 2,6-di-*alkyl*-pyridines such as 2,6-lutidine or 2,6-di-*tert*-butylpyridine. Suitable solvents include dimethylacetamide, dichloromethane, benzene, tetrahydrofuran and dimethylformamide. The coupling reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

Suitable activated acid derivatives include acid halides, for example acid chlorides, and active esters, for example pentafluorophenyl esters. The reaction of these types of compounds with amines is well known in the art, for example they may be reacted in the presence of a base, such as those described above, and in a suitable solvent, such as those
 5 described above. The reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

Compounds of formula (VI) may be prepared according to the following scheme:




wherein Pg is a suitable acid protecting group such as those described hereinbelow.

10 Amines of formula (VII) are commercially available or are prepared by processes known in the art.

Process d)

Acids of formula (IX) and amines of formula (VIII) may be coupled together under the conditions described in process c) above.

15 Amines of formula (VIII) may be prepared according to the following scheme:



wherein Pg is a suitable amino protecting group such as those described hereinbelow.

Acids of formula (IX) are commercially available or are prepared by processes known in the art.


20 Process e)

Acids of formula (VI) and amines of formula (X) may be coupled together under the conditions described in process c) above.

Amines of formula (X) are commercially available or are prepared by processes known in the art.


5 Examples of conversions of a compound of formula (I) into another compound of formula (I) are:

i) where G is $-NR^2-$; conversion of R^2 as hydrogen into other R^2 for example:



wherein L is a displaceable group;

10 ii) where G is $-NR^2-$; conversion of R^2 as a substituted side chain into another substituted side chain, for example:




wherein Ms is methanesulphonyl, and Nu is a nucleophile that introduces a substituent that is an optional substituent for R^2 as defined in formula (I) (NB the hydroxyl moiety does not

15 necessarily have to be on the terminal carbon as depicted above);

iii) conversion of one side chain of formula (Ia) into another side chain of formula (Ia).

iv) conversion of one value of R^1 into another value of R^1 , using standard techniques, for example, conversion of R^1 as hydroxy into C_{1-4} alkoxy.

The skilled reader will appreciate that the formation of the side chain (Ia) or (Ia') described in Processes c), d) and e) above and of the sidechain R² in i) and ii) above may also be performed on intermediates. For example:



5 A preferred process of the invention is Process b).

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of

10 the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the

15 introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halo group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with

20 a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where

protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Green, Protective Groups in Organic Synthesis, John Wiley and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may
5 be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection
10 conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid
15 as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by
20 treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl
25 group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group,
30 for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic

acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

- 5 Many of the intermediates defined herein are novel, for example, those of the formula II and IV and these are provided as a further feature of the invention.

ASSAYS

- As stated hereinbefore the pyrimidine derivative defined in the present invention possesses anti-cell-proliferation activity such as anti-cancer activity which is believed to arise
10 from the CDK and/or FAK inhibitory activity of the compound. These properties may be assessed, for example, using the procedure set out below:-

CDK4 Inhibition Assay

The following abbreviations have been used :-

HEPES is *N*-(2-Hydroxyethyl)piperazine-*N'*-(2-ethanesulfonic acid)

- 15 DTT is Dithiothreitol

PMSF is Phenylmethylsulfonyl fluoride

- The compounds were tested in an *in vitro* kinase assay in 96 well format using Scintillation Proximity Assay (SPA - obtained from Amersham) for measuring incorporation of [γ -33-P]-Adenosine Triphosphate into a test substrate (GST-Retinoblastoma). In each well
20 was placed the compound to be tested (diluted in DMSO and water to correct concentrations) and in control wells either p16 as an inhibitor control or DMSO as a positive control.

- Approximately 0.5 μ l of CDK4/Cyclin D1 partially-purified enzyme (amount dependent on enzyme activity) diluted in 25 μ l incubation buffer was added to each well then 20 μ l of GST-Rb/ATP/ATP33 mixture (containing 0.5 μ g GST-Rb and 0.2 μ M ATP and
25 0.14 μ Ci [γ -33-P]-Adenosine Triphosphate), and the resulting mixture shaken gently, then incubated at room temperature for 60 minutes.

- To each well was then added 150 μ L stop solution containing (0.8mg/well of Protein A-PVT SPA bead (Amersham)), 20pM/well of Anti-Glutathione Transferase, Rabbit IgG (obtained from Molecular Probes), 61mM EDTA and 50mM HEPES pH 7.5 containing
30 0.05% sodium azide.

The plates were sealed with Topseal-S plate sealers, left for two hours then spun at 2500rpm, 1124xg., for 5 minutes. The plates were read on a Topcount for 30 seconds per well.

The incubation buffer used to dilute the enzyme and substrate mixes contained 50mM HEPES pH7.5, 10mM MnCl₂, 1mM DTT, 100µM Sodium vanadate, 100µM NaF, 10mM Sodium Glycerophosphate, BSA (1mg/ml final).

As a control, another known inhibitor of CDK4 may be used in place of p16.

5 Test substrate

In this assay only part of the retinoblastoma (Science 1987 Mar13;235(4794):1394-1399; Lee W.H., Bookstein R., Hong F., Young L.J., Shew J.Y., Lee E.Y.) was used, fused to a GST tag. PCR of retinoblastoma amino acids 379-928 (obtained from retinoblastoma plasmid ATCC pLRbRNL) was performed, and the sequence cloned into
10 pGEX 2T fusion vector (Smith D.B. and Johnson, K.S. Gene 67, 31 (1988); which contained a tac promoter for inducible expression, internal lac I^q gene for use in any E.Coli host, and a coding region for thrombin cleavage - obtained from Pharmacia Biotech) which was used to amplify amino acids 792-928. This sequence was again cloned into pGEX 2T.

The retinoblastoma 792-928 sequence so obtained was expressed in E.Coli (BL21
15 (DE3) pLysS cells) using standard inducible expression techniques, and purified as follows.

E.coli paste was resuspended in 10ml/g of NETN buffer (50mM Tris pH 7.5, 120mM NaCl, 1mM EDTA, 0.5%v/v NP-40, 1mM PMSF, 1ug/ml leupeptin, 1ug/ml aprotinin and 1ug/ml pepstatin) and sonicated for 2 x 45 seconds per 100ml homogenate. After centrifugation, the supernatant was loaded onto a 10ml glutathione Sepharose column
20 (Pharmacia Biotech, Herts, UK), and washed with NETN buffer. After washing with kinase buffer (50mM HEPES pH 7.5, 10mM MgCl₂, 1mM DTT, 1mM PMSF, 1ug/ml leupeptin, 1ug/ml aprotinin and 1ug/ml pepstatin) the protein was eluted with 50mM reduced glutathione in kinase buffer. Fractions containing GST-Rb(792-927) were pooled and dialysed overnight against kinase buffer. The final product was analysed by Sodium Dodeca Sulfate
25 (SDS) PAGE (Polyacrylamide gel) using 8-16% Tris-Glycine gels (Novex, San Diego, USA).
CDK4 and Cyclin D1

CDK4 and Cyclin D1 were cloned from RNA from MCF-7 cell line (obtained from ATCC number:HTB22, breast adenocarcinoma line) as follows. The RNA was prepared from MCF-7 cells, then reverse transcribed using oligo dT primers. PCR was used to amplify the
30 complete coding sequence of each gene [CDK4 amino acids 1-303; Ref. Cell 1992 Oct 16; 71(2): 323-334; Matsushima H., Ewen M.E., Stron D.K., Kato J.Y., Hanks S.K., Roussel M.F., Sherr C.J. and Cyclin D1 amino acids 1-296; Ref. Cold Spring Harb. Symp. Quant.

Biol., 1991; 56:93-97; Arnold A., Motokura T., Bloom T., Kronenburg, Ruderman J., Juppner H., Kim H.G.].

After sequencing the PCR products were cloned using standard techniques into the insect expression vector pVL1393 (obtained from Invitrogen 1995 catalogue number :

- 5 V1392-20). The PCR products were then dually expressed [using a standard virus Baculogold co-infection technique] into the insect SF21 cell system (Spodoptera Frugiperda cells derived from ovarian tissue of the Fall Army Worm -Commercially available).

The following Example provides details of the production of Cyclin D1/CDK4 in SF21 cells (in TC100 + 10% FBS(TCS) + 0.2% Pluronic) having dual infection MOI 3 for
10 each virus of Cyclin D1 & CDK4.

Example production of Cyclin D1/CDK4

SF21 cells grown in a roller bottle culture to 2.33×10^6 cells/ml were used to inoculate 10 x 500 ml roller bottles at 0.2×10^6 cells/ml. The roller bottles were incubated on a roller rig at 28°C.

- 15 After 3 days (72 hrs.) the cells were counted, and the average from 2 bottles found to be 1.86×10^6 cells/ml. (99% viable). The cultures were then infected with the dual viruses at an MOI 3 for each virus.

10 x 500ml were infected with JS303 Cyclin D1 virus titre - 9×10^7 pfu/ml. JS304 CDK4 virus titre - 1×10^8 pfu/ml.

- 20 Cyclin D1 $\frac{1.86 \times 10^6 \times 500 \times 3}{0.9 \times 10^8} = 31$ ml of virus for each 500 ml. bottle.

CDK4 $\frac{1.86 \times 10^6 \times 500 \times 3}{1 \times 10^8} = 28$ ml of virus for each 500 ml. bottle.

- 25 The viruses were mixed together before addition to the cultures, and the cultures returned to the roller rig 28°C.

- After 3 days (72 hrs.) post infection the 5 Litres of culture was harvested. The total cell count at harvest was 1.58×10^6 cells/ml.(99% viable). The cells were spun out at 2500rpm, 30 mins., 4°C in Heraeus Omnifuge 2.0 RS in 250 ml lots. The supernatant was discarded. 20 pellets of $\sim 4 \times 10^8$ cells/pellet were snap frozen in LN₂ and stored at -80°C in
30 CCRF cold room. The SF21 cells were then hypotonically lysed by resuspending in lysis buffer (50mM HEPES pH 7.5, 10mM magnesium chloride, 1mM DTT, 10mM glycerophosphate, 0.1mM PMSF, 0.1mM sodium fluoride, 0.1mM sodium orthovanadate,

5 $\mu\text{g/ml}$ aprotinin, $5 \mu\text{g/ml}$ leupeptin and 20% w/v sucrose), and adding ice cold deionised water. After centrifugation, the supernatant was loaded onto a Poros HQ/M 1.4/100 anion exchange column (PE Biosystems, Hertford, UK). CDK4 and Cyclin D1 were coeluted with 375mM NaCl in lysis buffer, and their presence checked by western blot, using suitable anti-CDK4 and anti-Cyclin D1 antibodies (obtained from Santa Cruz Biotechnology, California, US).

p16 control (Nature 366:704-707; 1993; Serrano M, Hannon GJ, Beach D)

p16 (the natural inhibitor of CDK4/Cyclin D1) was amplified from HeLa cDNA (Hela cells obtained from ATCC CCL2, human epitheloid carcinoma from cervix; Cancer Res. 12: 10 264, 1952), cloned into pTB 375 NBSE which had a 5' His tag, and transformed using standard techniques into BL21 (DE3) pLysS cells (obtained from Promega; Ref. Studier F.W. and Moffat B.A., J. Mol. Biol., 189, 113, 1986). A 1 litre culture was grown to the appropriate OD then induced with IPTG to express p16 overnight. The cells were then lysed by sonication in 50mM sodium phosphate, 0.5M sodium chloride, PMSF, $0.5 \mu\text{g/ml}$ leupeptin and $0.5 \mu\text{g/ml}$ 15 aprotinin. The mixture was spun down, the supernatant added to nickel chelate beads and mixed for 1 ½ hours. The beads were washed in sodium phosphate, NaCl pH 6.0 and p16 product eluted in sodium phosphate, NaCl pH 7.4 with 200mM imidazole.

The pTB NBSE was constructed from pTB 375 NBPE as follows :-

p TB375

20 The background vector used for generation of pTB 375 was pZEN0042 (see UK patent 2253852) and contained the tetA/tetR inducible tetracycline resistance sequence from plasmid RP4 and the cer stability sequence from plasmid pKS492 in a pAT153 derived background. pTB375 was generated by the addition of an expression cassette consisting of the T7 gene 10 promoter, multiple cloning site and T7 gene 10 termination sequence. In addition, a terminator 25 sequence designed to reduce transcriptional readthrough from the background vector was included upstream of the expression cassette.

pTB 375 NBPE

The unique EcoRI restriction site present in pTB 375 was removed. A new multiple cloning site containing the recognition sequences for the restriction enzymes NdeI, BamHI, 30 PstI and EcoRI was introduced into pTB 375 between the NdeI and BamHI sites destroying the original BamHI site present in pTB 375.

pTB 375 NBSE

A new multiple cloning site containing the recognition sequences for the restriction enzymes NdeI, BamHI, SmaI and EcoRI was introduced into pTB 375 NBPE between the NdeI and EcoRI sites. The oligonucleotide containing these restriction sites also contained 6
5 histidine codons located between the NdeI and BamHI sites in the same reading frame as the initiator codon (ATG) present within the NdeI site.

By analogy to the above, assays designed to assess inhibition of CDK2 and CDK6 may be constructed. CDK2 (EMBL Accession No. X62071) may be used together with Cyclin A or Cyclin E (see EMBL Accession No. M73812), and further details for such assays
10 are contained in PCT International Publication No. WO99/21845, the relevant Biochemical & Biological Evaluation sections of which are hereby incorporated by reference.

If using CDK2 with Cyclin E partial co-purification may be achieved as follows:- Sf21 cells are resuspended in lysis buffer (50mM Tris pH 8.2, 10mM MgCl₂, 1mM DTT, 10mM glycerophosphate, 0.1mM sodium orthovanadate, 0.1mM NaF, 1mM PMSF, 1ug/ml
15 leupeptin and 1ug/ml aprotinin) and homogenised for 2 minutes in a 10ml Dounce homogeniser. After centrifugation, the supernatant is loaded onto a Poros HQ/M 1.4/100 anion exchange column (PE Biosystems, Hertford, UK). CDK2 and Cyclin E are coeluted at the beginning of a 0-1M NaCl gradient (run in lysis buffer minus protease inhibitors) over 20 column volumes. Co-elution is checked by western blot using both anti-CDK2 and
20 anti-Cyclin E antibodies (Santa Cruz Biotechnology, California, US).

FAK3 Kinase Inhibition Assay

This assay determines the ability of a test compound to inhibit tyrosine kinase activity of human Focal Adhesion Kinase (FAK).

DNA encoding FAK is obtained by total gene synthesis (Edwards M, International
25 Biotechnology Lab 5(3), 19-25, 1987) or by cloning. These are then expressed in a suitable expression system to obtain polypeptide with tyrosine kinase activity. For example, FAK, obtained by expression of recombinant protein in insect cells, was found to display intrinsic tyrosine kinase activity.

FAK (full length human cDNA described by Andre et al (Biochemical and Biophysical
30 Research Communications, 1993, 190 (1): 140-147; EMBL/GenBank Accession Number L05186)) was modified such that the resulting protein when translated had a 6-histidine tag at the N-terminus immediately preceding the start methionine. Active FAK protein has been

previously expressed in a baculovirus system using a similar N-terminal 6-histidine tag (Protein Expression And Purification, 1996, 7: 12-18). The human FAK cDNA was cloned into the baculovirus transplacement vector, pFastbac 1 (Life Technologies), and the recombinant construct was co-transfected into insect cells (for example *Spodoptera frugiperda* 21(Sf21))

- 5 with viral DNA to prepare recombinant baculovirus (details of the methods for the assembly of recombinant DNA molecules and the preparation and use of recombinant baculovirus can be found in standard texts for example Sambrook et al, 1989, Molecular cloning - A Laboratory Manual, 2nd edition, Cold Spring Harbour Laboratory Press and O'Reilly et al, 1992, Baculovirus Expression Vectors - A Laboratory Manual, W. H. Freeman and Co, New York.
- 10 Details specific to the use of the pFastbac ('Bac to Bac') system are provided in Anderson et al., 1995, FOCUS (Life Technologies Bulletin Magazine), 17, p53.)

- For expression of biologically active human FAK protein, Sf21 cells were infected with plaque-pure FAK recombinant virus at a multiplicity of infection of 3 and harvested 48 hours later. Harvested cells were washed with ice cold phosphate buffered saline solution (PBS)
- 15 (10mM sodium phosphate pH7.4, 138mM sodium chloride, 2.7mM potassium chloride) then resuspended in ice cold lysis buffer (50mM HEPES pH7.5, 1mM Dithiothreitol, 100uM Sodium Fluoride, 100uM Sodium Orthovanadate, 10mM Glycerophosphate, 100uM Phenylmethylsulphonylfluoride (PMSF), 5ug/ml Aprotinin, 5ug/ml Leupeptin, 1% Tween; the PMSF being added just before use from a freshly-prepared 100mM solution in methanol) using
- 20 250µl lysis buffer per 10 million cells. The suspension was then incubated on ice for 15 minutes and centrifuged for 10 minutes at 13,000 rpm at 4°C. The supernatant (enzyme stock) was removed and aliquots made which were snap frozen in liquid nitrogen and then stored at -70°C. For a typical batch, stock enzyme was diluted 1 in 250 with enzyme diluent ((100mM HEPES pH 7.4, 0.2mM Dithiothreitol, 200uM Sodium Orthovanadate, 0.1% Triton X-100) and
- 25 50ml of freshly diluted enzyme was used for each assay well (see FAK3 protocol, below).

FAK3: *In vitro* Enzyme assay Protocol

A stock of substrate solution was prepared from a random copolymer containing tyrosine, for example Poly (Glu, Ala, Tyr) 6:3:1 (Sigma P3899), stored as 1 mg/ml stock in PBS at -20°C and diluted 1 in 500 with PBS for plate coating.

- 30 On the day before the assay 100µl of diluted substrate solution was dispensed into all wells of assay plates (Maxisorp 96 well immunoplates Life technologies, Cat. No. 439454A) which were sealed with plate sealers and left overnight at 4°C.

On the day of the assay the substrate solution was discarded and the assay plate wells were washed once with 200 μ l PBST (PBS containing 0.05% v/v Tween 20) and once with 200 μ l 50mM Hepes pH7.4.

Test compounds were made up as 10mM or 30mM stocks in DMSO and then further
5 diluted in glass distilled water diluted to a concentration 10 fold higher than the final assay concentration. 10 μ l of diluted compound was transferred to wells in the washed assay plates. "No compound" control wells contained 10 μ l glass distilled water instead of compound.

Forty microlitres of 25mM manganese chloride containing 6.25 μ M
adenosine-5'-triphosphate (ATP) was added to all test wells. To start the reactions 50 μ l of
10 freshly diluted enzyme was added to each well and the plates were incubated at 23C for 90 minutes. Then the reaction was stopped by adding 100 μ l of PBS containing 20mM EDTA. The liquid was then discarded and the wells were washed twice with PBST.

One hundred microlitres of mouse HRP-linked anti-phosphotyrosine antibody (Santa
Cruz, Product SC 7020-HRP), diluted 1 in 1500 with PBST containing 0.5% w/v bovine serum
15 albumin (BSA), was added to each well and the plates were incubated for 1 hour at room temperature before discarding the liquid and washing the wells twice with 200 μ l PBST. One hundred microlitres of 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulphonic acid) (ABTS) solution, freshly prepared using one 50mg ABTS tablet (Boehringer 1204 521) in 50ml freshly prepared 50mM phosphate-citrate buffer pH5.0 + 0.03% sodium perborate (made with 1 phosphate
20 citrate buffer with sodium perborate (PCSB) capsule (Sigma P4922) per 100ml distilled water), was added to each well. Plates were then incubated for 20-60 minutes at room temperature until the absorbance value of the "no compound" control wells, measured at 405nm using a plate reading spectrophotometer, was approximately 1.0.

Dose response curves were generated from the absorbance readings using Origin
25 Software. Compounds were ranked for potency using the Inhibitory Concentration 50 (IC₅₀), as defined by Origin Software analysis.

Although the pharmacological properties of the compounds of the formula (I) vary
with structural change, in general activity possessed by compounds of the formula (I) in the
above assays may be demonstrated at IC₅₀ concentrations or doses in the range 250 μ M to
30 1nM.

When tested in the above *in vitro* assay the CDK4 inhibitory activity of Example 45
was measured as IC₅₀ = 0.235 μ M. When tested in the above *in vitro* assay the FAK inhibitory

activity of Example 47 was measured as $IC_{50} = 0.097\mu M$ and that of Example 52 as $IC_{50} = 0.814\mu M$.

The *in vivo* activity of the compounds of the present invention may be assessed by standard techniques, for example by measuring inhibition of cell growth and assessing

5 cytotoxicity. For example, further details may be found in the following references:-

a) Attenuation of the Expression of the Focal Adhesion Kinase induces Apoptosis in Tumor Cells. Xu L-h et al. Cell Growth & Differentiation (1996) 7, p413-418;

b) The COOH-Terminal Domain of the Focal Adhesion Kinase Induces Loss of Adhesion and Cell Death in Human Tumour Cells. Xu L-h et al. Cell Growth & Differentiation (1998) 9,

10 p999-1005;

c) Inhibition of pp125-FAK in Cultured Fibroblasts Results in Apoptosis. Hungerford J.E et al. The Journal of Cell Biology (1996) 135, p1383-1390;

d) Inhibition of Focal Adhesion Kinase (FAK) Signalling in Focal Adhesions Decreases Cell Motility and Proliferation. Gilmore A.P and Romer L.H. Molecular Biology of the Cell

15 (1996) 7, p1209-1224.

Inhibition of cell growth may be measured by staining cells with Sulforhodamine B (SRB), a fluorescent dye that stains proteins and therefore gives an estimation of amount of protein (i.e. cells) in a well (see Boyd, M. R. (1989) Status of the NCI preclinical antitumour drug discovery screen. Prin. Prac Oncol 10:1-12). Thus, the following details are provided of

20 measuring inhibition of cell growth:-

Cells were plated in appropriate medium in a volume of 100 μ l in 96 well plates; media was Dulbecco's Modified Eagle media for MCF-7, SK-UT-1B and SK-UT-1. The cells were allowed to attach overnight, then inhibitor compounds were added at various concentrations in a maximum concentration of 1% DMSO (v/v). A control plate was assayed to give a value for

25 cells before dosing. Cells were incubated at 37°C, (5% CO₂) for three days.

At the end of three days TCA was added to the plates to a final concentration of 16% (v/v). Plates were then incubated at 4°C for 1 hour, the supernatant removed and the plates washed in tap water. After drying, 100 μ l SRB dye (0.4% SRB in 1% acetic acid) was added for 30 minutes at 37°C. Excess SRB was removed and the plates washed in 1% acetic acid.

30 The SRB bound to protein was solubilised in 10mM Tris pH7.5 and shaken for 30 minutes at room temperature. The ODs were read at 540nm, and the concentration of inhibitor causing 50% inhibition of growth was determined from a semi-log plot of inhibitor concentration

versus absorbance. The concentration of compound that reduced the optical density to below that obtained when the cells were plated at the start of the experiment gave the value for toxicity.

Typical IC_{50} values for compounds of the invention when tested in the SRB assay are
5 in the range 1mM to 1nM.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in association with a pharmaceutically acceptable diluent or carrier.

10 The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository.

In general the above compositions may be prepared in a conventional manner using
15 conventional excipients.

The pyrimidine will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000 mg per square meter body area of the animal, i.e. approximately 0.1-100 mg/kg, and this normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain, for example 1-250 mg of active ingredient.
20 Preferably a daily dose in the range of 1-50 mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

According to a further aspect of the present invention there is provided a pyrimidine
25 derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

We have found that the pyrimidine derivatives defined in the present invention, or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, are effective cell cycle
30 inhibitors (anti-cell proliferation agents), which property (without being bound by theory) is believed to arise from their CDK inhibitory properties. The compounds are also effective inhibitors of FAK. Accordingly the compounds of the present invention are expected to be

useful in the treatment of diseases or medical conditions mediated alone or in part by CDK and/or FAK enzymes, i.e. the compounds may be used to produce a CDK and/or FAK inhibitory effect in a warm-blooded animal in need of such treatment. Thus the compounds of the present invention provide a method for treating the proliferation and/or migration of

5 malignant cells characterised by inhibition of CDK and/or FAK enzymes, i.e. the compounds may be used to produce an anti-proliferative/migration effect mediated alone or in part by the inhibition of CDKs and/or FAK. The compounds may also be useful as FAK inhibitors by inducing cell-death (apoptosis). Such a pyrimidine derivative of the invention is expected to possess a wide range of anti-cancer properties as CDKs and/or FAK have been implicated in

10 many common human cancers such as leukaemia and breast, lung, colon, rectal, stomach, prostate, bladder, pancreas and ovarian cancer. Thus it is expected that a pyrimidine derivative of the invention will possess anti-cancer activity against these cancers. It is in addition expected that a pyrimidine derivative of the present invention will possess activity against a range of leukaemias, lymphoid malignancies and solid tumours such as carcinomas and

15 sarcomas in tissues such as the liver, kidney, prostate and pancreas. In particular such compounds of the invention are expected to slow advantageously the growth of primary and recurrent solid tumours of, for example, the colon, breast, prostate, lungs and skin. More particularly such compounds of the invention, or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, are expected to inhibit the growth of those primary and recurrent

20 solid tumours which are associated with CDK and/or FAK, especially those tumours which are significantly dependent on CDK and/or FAK for their growth and spread, including for example, certain tumours of the colon, breast, prostate, lung, vulva and skin.

It is further expected that a pyrimidine derivative of the present invention will possess activity against other cell-proliferation/migration diseases in a wide range of other disease

25 states including leukemias, fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic inflammation, bone diseases and ocular diseases with retinal vessel proliferation.

Thus according to this aspect of the invention there is provided a pyrimidine derivative

30 of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore for use as a medicament; and the use of a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as

defined hereinbefore in the manufacture of a medicament for use in the production of an anti-cancer, cell cycle inhibitory (anti-cell-proliferation) effect and/or a FAK inhibitory (anti-cell migration and/or apoptosis inducing) effect in a warm-blooded animal such as man. Particularly, a cell cycle inhibitory effect is produced at the S or G1-S phase by inhibition of

5 CDK2, CDK4 and/or CDK6, especially CDK4 and CDK6.

According to a further feature of this aspect of the invention there is provided a method for producing an anti-cancer, cell cycle inhibitory (anti-cell-proliferation) effect and/or a FAK inhibitory (anti-cell migration and/or apoptosis inducing) effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering

10 to said animal an effective amount of a pyrimidine derivative as defined immediately above. Particularly, an inhibitory effect is produced at the S or G1-S phase by inhibition of CDK2, CDK4 and/or CDK6, especially CDK4 and CDK6.

As stated above the size of the dose required for the therapeutic or prophylactic treatment of a particular cell-proliferation disease will necessarily be varied depending on the

15 host treated, the route of administration and the severity of the illness being treated. A unit dose in the range, for example, 1-100 mg/kg, preferably 1-50 mg/kg is envisaged.

The CDK and/or FAK inhibitory activity defined hereinbefore may be applied as a sole therapy or may involve, in addition to a compound of the invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the

20 simultaneous, sequential or separate administration of the individual components of the treatment. In the field of medical oncology it is normal practice to use a combination of different forms of treatment to treat each patient with cancer. In medical oncology the other component(s) of such conjoint treatment in addition to the cell cycle inhibitory treatment defined hereinbefore may be: surgery, radiotherapy or chemotherapy. Such chemotherapy

25 may cover three main categories of therapeutic agent:

- (i) other cell cycle inhibitory agents that work by the same or different mechanisms from those defined hereinbefore;
- (ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene, idoxifene), progestogens (for example megestrol acetate), aromatase inhibitors

30 (for example anastrozole, letrozole, vorazole, exemestane), antiprogestogens, antiandrogens (for example flutamide, nilutamide, bicalutamide, cyproterone acetate), LHRH agonists and antagonists (for example goserelin acetate, luprolide), inhibitors of testosterone

5 α -dihydroreductase (for example finasteride), anti-invasion agents (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator receptor function) and inhibitors of growth factor function, (such growth factors include for example platelet derived growth factor and hepatocyte growth factor such inhibitors include growth factor antibodies, growth factor receptor antibodies, tyrosine kinase inhibitors and serine/threonine kinase inhibitors); and

(iii) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as antimetabolites (for example antifolates like methotrexate, fluoropyrimidines like 5-fluorouracil, purine and adenosine analogues, cytosine arabinoside);

antitumour antibiotics (for example anthracyclines like doxorubicin, daunomycin, epirubicin and idarubicin, mitomycin-C, dactinomycin, mithramycin); platinum derivatives (for example cisplatin, carboplatin); alkylating agents (for example nitrogen mustard, melphalan, chlorambucil, busulphan, cyclophosphamide, ifosfamide, nitrosoureas, thiotepa); antimetotic agents (for example vinca alkaloids like vincristine and taxoids like taxol, taxotere);

topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan). According to this aspect of the invention there is provided a pharmaceutical product comprising a pyrimidine derivative of the formula (I) as defined hereinbefore, or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, and an additional anti-tumour substance as defined hereinbefore for the conjoint treatment of cancer. An anti-emetic may also be usefully administered, for example when using such conjoint treatment as described above.

In addition to their use in therapeutic medicine, the compounds of formula (I) and their pharmaceutically acceptable salts are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of cell cycle activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

In the above other, pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

The invention will now be illustrated in the following non limiting Examples, in which standard techniques known to the skilled chemist and techniques analogous to those described in these Examples may be used where appropriate, and in which, unless otherwise stated:

- (i) evaporations were carried out by rotary evaporation *in vacuo* and work up procedures were carried out after removal of residual solids such as drying agents by filtration;
- (ii) operations were carried out at ambient temperature, typically in the range 18-25°C and in air unless stated, or unless the skilled person would otherwise operate under an atmosphere of
- 5 an inert gas such as argon;
- (iii) column chromatography (by the flash procedure) and medium pressure liquid chromatography (MPLC) were performed on Merck Kieselgel silica (Art. 9385) or on Merck Lichroprep RP-18 (Art. 9303) reversed-phase silica, obtained from E. Merck, Darmstadt, Germany; bond elute chromatography was performed using Varian Mega Bond Elut
- 10 cartridges (10 g, order code 1225-6034), obtained from Varian Sample Preparation Products, California, USA;
- (iv) yields are given for illustration only and are not necessarily the maximum attainable;
- (v) the structures of the end products of the formula (I) were generally confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; proton magnetic
- 15 resonance chemical shift values were measured in deuterated DMSO-d₆ (unless otherwise stated) on the delta scale (ppm downfield from tetramethylsilane) using a Varian Gemini 2000 spectrometer operating at a field strength of 300MHz, or a Bruker AM250 spectrometer operating at a field strength of 250MHz; and peak multiplicities are shown as follows: s, singlet; d, doublet; dd, double doublet; t, triplet; tt, triple triplet; q, quartet; tq, triple quartet;
- 20 m, multiplet; br, broad; mass spectrometry (MS) was performed by electrospray on a VG platform;
- (vi) unless further details are specified in the text, analytical high performance liquid chromatography (HPLC) was performed on a Waters Spherisorb ODS1 25 cm column, at a flow rate of 2 ml/minute using acetonitrile/water/trifluoroacetic acid (60:40:0.1 v/v) as eluent,
- 25 detection was at a wavelength of 254 nm, and data are quoted as retention time (RT) in minutes;
- (vii) robotic synthesis was carried out using a Zymate XP robot, with solution additions via a Zymate Master Laboratory Station and stirred via a Stem RS5000 Reacto-Station at 25°C;
- (viii) work up and purification of reaction mixtures from robotic synthesis was carried out as
- 30 follows: evaporations were carried out *in vacuo* using a Savant AES 2000; column chromatography was performed using either an Anachem Sympur MPLC or Jones Flashmaster MPLC systems on silica using Varian Mega Bond Elut cartridges; the structures

of the final products were confirmed by LCMS on a Micromass OpenLynx system using the following and are quoted as retention time (RT) in minutes:

Column: 4.6 mm x 10 cm Hichrom RPB 100Å

Solvent (System A): I = 5% Methanol in Water + 0.1% formic acid,

5 II = 5% Methanol in Acetonitrile + 0.1% formic acid

Solvent (System B): I = Water + 0.1% formic acid,

II = Acetonitrile + 0.1% formic acid

Run time: 10 minutes with a 6 minute gradient from 5-95% II

Wavelength: 254 nm, bandwidth 10 nm

10 Mass detector: Platform LC

(ix) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), HPLC, infra-red (IR), MS or NMR analysis;

(x) where solutions are dried magnesium sulphate was the drying agent;

(xi) the following abbreviations may be used hereinbefore or hereinafter:-

15 DCM dichloromethane;
DMF *N,N*-dimethylformamide;
DMSO dimethylsulphoxide;
NMP *N*-methylpyrrolidin-2-one;
THF tetrahydrofuran; and

20 (xii) for the avoidance of doubt, where "precursor amines" are described and this amine contains more than one nitrogen, the resulting example formed is not a quaternary compound.

Example 1

4-Anilino-5-chloro-2-{4-{*N*-[3-(imidazol-1-yl)propyl]carbamoyl}anilino}pyrimidine

25 4-Anilino-2-(4-carboxyanilino)-5-chloropyrimidine (Method 1; 170 mg, 0.5 mmol) was added to 1-(3-aminopropyl)imidazole (93.9 mg, 0.75 mmol) in a reaction tube.

1-Hydroxybenzotriazole (101 mg, 0.75 mmol) was added followed by

N,N-diisopropylethylamine (130 ml, 0.75 mmol) in DCM (4 ml). The mixture was flushed with argon and then stirred for 1 hour. 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide


30 hydrochloride (105 mg, 0.55 mmol) in DCM (3 ml) was added and the mixture was stirred for 16 hours. The mixture was washed with saturated sodium bicarbonate solution (10 ml), water (10 ml) and saturated sodium chloride solution (10 ml), and the organic phase was loaded on a

Varian Mega Bond Elut column. Elution with 0-10% 2.0M methanolic ammonia solution in DCM gave the product (11.6 mg, 5.2%). NMR: 2.0-2.15 (m, 2H), 3.0-3.1 (m, 2H), 4.25 (t, 2H), 7.25 (t, 1H), 7.4 (t, 2H), 7.6 (d, 4H), 7.7 (s, 1H), 7.75 (d, 2H), 7.85 (d, 1H), 8.3 (s, 1H), 8.6 (br t, 1H), 9.25 (d, 1H), 9.6 (br s, 1H), 10.3 (br s, 1H); LCMS (MH⁺): 448; HPLC (RT, System A): 6.16.

Examples 2-10

The following compounds were prepared using a Zymate XP robot by an analogous method to that described in Example 1, using

- 10 4-anilino-2-(4-carboxyanilino)-5-chloropyrimidine (Method 1) and the appropriate amine:



Ex	R (Precursor amine)	LCMS (MH ⁺)	HPLC ¹ (RT)
2	1-(3-Aminopropyl)-2-pyrrolidinone	465	6.5
3	4-(2-Aminoethyl)morpholine	453	5.99
4	4-(3-Aminopropyl)morpholine	467	6.04
5	1-(2-Aminoethyl)pyrrolidine	437	6.12
6	2-(Diethylamino)ethylamine	439	6.09
7	1-(3-Aminopropyl)-4-methylpiperazine	480	5.7
8	1-(3-Aminopropyl)pyrrolidine	451	5.95
9	3-(2-Aminoethyl)-3-aza-8-oxabicyclo[3,2,1]hexane ²	479	6.42
10	4-(2-Aminoethyl)thiomorpholine ³	469	6.27

¹ System A


² Precursor amine obtained as described in US Pat. Appl. 3856783

³ Precursor amine obtained as described in J. Med. Chem. (1974), 17, 1232-4

Examples 11-20

The following compounds were prepared using a Zymate XP robot by an analogous method to that described in Example 1, using

- 5 4-anilino-5-bromo-2-(3-carboxyanilino)pyrimidine (Method 4) and the appropriate amine:



Ex	R (Precursor amine)	LCMS (MH ⁺)	HPLC ¹ (RT)
11	1-(3-Aminopropyl)pyrrolidine	495	5.94
12	3-(2-Aminoethyl)-3-aza-8-oxabicyclo[3,2,1]hexane ²	523	6.25
13	4-(2-Aminoethyl)thiomorpholine ³	513	6.25
14	1-(3-Aminopropyl)imidazole	492	5.8
15	1-(3-Aminopropyl)-2-pyrrolidinone	509	6.4
16	4-(2-Aminoethyl)morpholine	497	5.86
17	4-(3-Aminopropyl)morpholine	511	6.36
18	1-(2-Aminoethyl)pyrrolidine	481	5.99
19	2-(Diethylamino)ethylamine	483	6.05
20	1-(3-Aminopropyl)-4-methylpiperazine	524	5.68

¹ System A

² Precursor amine obtained as described in US Pat. Appl. 3856783

³ Precursor amine obtained as described in J. Med. Chem. (1974), 17, 1232-4


10

Examples 21-30

The following compounds were prepared using a Zymate XP robot by an analogous method to that described in Example 1, using

5-bromo-2-(4-carboxyanilino)-4-(4-methoxyanilino)pyrimidine (Method 5) and the

- 15 appropriate amine:



Ex	R (Precursor amine)	LCMS (MH ⁺)	HPLC ¹ (RT)
21	1-(3-Aminopropyl)imidazole	522	6.09
22	1-(3-Aminopropyl)-2-pyrrolidinone	539	6.53
23	4-(2-Aminoethyl)morpholine	527	6.05
24	4-(3-Aminopropyl)morpholine	541	60.1
25	1-(2-Aminoethyl)pyrrolidine	511	6.05
26	2-(Diethylamino)ethylamine	513	6.16
27	1-(3-Aminopropyl)-4-methylpiperazine	554	5.75
28	1-(3-Aminopropyl)pyrrolidine	525	6.01
29	3-(2-Aminoethyl)-3-aza-8-oxabicyclo[3,2,1]hexane ²	553	6.43
30	4-(2-Aminoethyl)thiomorpholine ³	543	6.32

¹ System A

² Precursor amine obtained as described in US Pat. Appl. 3856783

³ Precursor amine obtained as described in J. Med. Chem. (1974), 17, 1232-4


5

Examples 31-40

The following compounds were prepared using a Zymate XP robot by an analogous method to that described in Example 1, using

5-bromo-2-(4-carboxyanilino)-4-(4-methoxyphenoxy) pyrimidine (Method 26) and the

10 appropriate amine:



Ex	R (Precursor amine)	LCMS (MH ⁺)	HPLC ¹ (RT)
31	1-(3-Aminopropyl)imidazole	523	6.82
32	1-(3-Aminopropyl)-2-pyrrolidinone	540	7.19
33	4-(2-Aminoethyl)morpholine	528	6.75
34	4-(3-Aminopropyl)morpholine	542	6.45
35	1-(2-Aminoethyl)pyrrolidine	512	6.79
36	2-(Diethylamino)ethylamine	514	6.86
37	1-(3-Aminopropyl)-4-methylpiperazine	555	6.55
38	1-(3-Aminopropyl)pyrrolidine	526	6.97
39	3-(2-Aminoethyl)-3-aza-8-oxabicyclo[3,2,1]hexane ²	554	6.91
40	4-(2-Aminoethyl)thiomorpholine ³	544	6.95


¹ System A² Precursor amine obtained as described in US Pat. Appl. 3856783³ Precursor amine obtained as described in J. Med. Chem. (1974), 17, 1232-4

5

Examples 41-58

The following compounds were prepared using a Zymate XP robot by an analogous method to that described in Example 1, using

4-anilino-5-bromo-2-(4-carboxyanilino)pyrimidine (Method 6) and the appropriate amine:



Ex	R (Precursor amine)	LCMS (MH ⁺)	HPLC ¹ (RT)
41	1-(3-Aminopropyl)imidazole	492	5.9
42	N-(3-Aminopropyl)-2-pyrrolidinone	509	6.3
43	Tryptamine	527	6.7
44	4-(3-Aminopropyl)morpholine	511	5.9
45	1-(2-Aminoethyl)piperidine	495	6.1
46	1-(2-Aminoethyl)pyrrolidine	481	6.0
47	3-(Di-n-butylamino)propylamine	553	6.4
48	3-(n-Butoxy)propylamine	498	7.0
49	3-(Methylthio)propylamine	472	6.6
50	2-(Di-n-butylamino)ethylamine	539	6.5
51	2-(Thien-2-yl)ethylamine	494	6.8
52	4-(Pyrrolidin-1-yl)piperidine	521	5.9
53	2-Aminomethylimidazo[1,2-a]pyridine	514	6.1
54	4-(2-Aminoethyl)morpholine	497	5.9
55	1-Aminoethyl-3,5-dimethylpyrazole	506	6.5
56	2-(Methylthio)ethylamine	458	7.6
57	1-(2-Aminoethyl)-2-methyl-5-nitroimidazole	537	6.4
58	2-Methylamine-5-methylfuran	478	6.7

¹ System B

² Precursor amine obtained as described in Heterocycl. Commun. (1996), 2, 241-246

³ Precursor amine obtained as described in Recl. Trav. Chim. Pays-Bas (1992), 111, 371-8

5 ⁴ Precursor amine obtained as described in Eur. J. Med. Chem. - Chim. Ther. (1975), 10,


171-7

Example 59**5-Bromo-4-(4-fluoroanilino)-2-(4-{N-[3-(2-oxopyrrolidin-1-yl)propyl]carbamoyl}anilino)pyrimidine**

1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (52 mg, 0.27 mmol) was added to a mixture of 5-bromo-2-(4-carboxyanilino)-4-(4-fluoroanilino)pyrimidine (Method 7; 100 mg, 0.25 mmol), *N*-(3-aminopropyl)-2-pyrrolidinone (38 mg, 0.27 mmol) and 1-hydroxybenzotriazole (44 mg, 0.32 mmol) in DMF (0.5 ml). The mixture was stirred for 16 hours and then a 1:1 mixture of saturated sodium chloride solution and water (10 ml) was added. The precipitated solid was collected by filtration and dried to a constant weight in a vacuum oven at 40°C to give the product as an off-white solid (98 mg, 75%). MS (MH⁺): 528; HPLC (RT): 4.01.

Examples 60-85

The following compounds were prepared by an analogous method to that described in Example 59, using the appropriate 4-anilino-5-bromo-2-(4-carboxyanilino)pyrimidine (Methods 7-13) and the appropriate amine:



Ex	R ¹	R ² (Precursor amine)	MS (MH ⁺)	HPLC(RT)
60	4-F	1-(2-Aminoethyl)pyrrolidine	499, 501	5.11
61	4-F	1-(2-Aminoethyl)piperidine	514, 516	5.80
62	4-F	2-Isopropylaminoethylamine ¹	488, 490	4.03
63	3,4-di-F	1-(3-Aminopropyl)-2-pyrrolidinone	545, 547	4.52
64	3,4-di-F	1-(2-Aminoethyl)pyrrolidine	518, 520	5.63
65	3,4-di-F	1-(2-Aminoethyl)piperidine	531, 533	7.09
66	3,4-di-F	2-Isopropylaminoethylamine ¹	505, 507	4.71


67	4-SMe	1-(3-Aminopropyl)-2-pyrrolidinone	555, 557	5.18
68	4-SMe	1-(2-Aminoethyl)pyrrolidine	527, 529	5.08
69	4-SMe	1-(2-Aminoethyl)piperidine	541, 543	7.12
70	4-SMe	2-Isopropylaminoethylamine ¹	516, 518	4.84
71	4-Br	1-(3-Aminopropyl)-2-pyrrolidinone	589, 591	5.72
72	4-Br	1-(2-Aminoethyl)pyrrolidine	561, 563	6.29
73	4-Br	1-(2-Aminoethyl)piperidine	575, 577	7.48
74	4-Br	2-Isopropylaminoethylamine ¹	549, 551	4.56
75	3-Me	1-(3-Aminopropyl)-2-pyrrolidinone	523, 525	5.10
76	3-Me	1-(2-Aminoethyl)pyrrolidine	495, 497	5.25
77	3-Me	1-(2-Aminoethyl)piperidine	509, 511	6.66
78	3-Me	2-Isopropylaminoethylamine ¹	483, 485	4.72
79	3-F	1-(3-Aminopropyl)-2-pyrrolidinone	527, 529	4.12
80	3-F	1-(2-Aminoethyl)pyrrolidine	499, 501	5.69
81	3-F	1-(2-Aminoethyl)piperidine	513, 515	6.58
82	3-F	2-Isopropylaminoethylamine ¹	487, 489	6.63
83	2-CH ₂ OH	1-(3-Aminopropyl)-2-pyrrolidinone	539, 541	2.53
84	2-CH ₂ OH	1-(2-Aminoethyl)pyrrolidine	511, 513	2.47
85	2-CH ₂ OH	2-Isopropylaminoethylamine ¹	499, 501	2.15

¹ Substituent in 4-position of the 2-anilino of the resulting example is

N-(2-isopropylaminoethyl)carbamoyl.

Examples 86-90


- 5 The following compounds were prepared by an analogous method to that described in Example 59, using 2-(4-carboxyanilino)-5-fluoro-4-(4-methoxyanilino)pyrimidine (Method 27) and the appropriate amine:



Ex	R (Precursor amine)	NMR	MS (MH ⁺)
86	4-(2-Aminoethyl) morpholine	2.31-2.48 (m, 6H), 3.22-3.40 (m, 2H), 3.81 (s, 3H), 7.05 (d, 2H), 7.25 (d, 2H), 7.47 (d, 2H), 7.55 (d, 2H), 8.08-8.17 (m, 1H), 8.50 (d, 1H), 9.78 (s, 1H)	468
87	1-(2-Aminoethyl) piperidine	1.27-1.48 (m, 6H), 2.23-2.47 (m, 6H), 3.23-3.42 (m, 2H), 3.82 (s, 3H), 7.05 (d, 2H), 7.24 (d, 2H), 7.46 (d, 2H), 7.54 (d, 2H), 8.10 (m, 1H), 8.49 (d, 1H), 9.78 (s, 1H)	466
88	4-(Dimethylamino)benzylamine	2.83 (s, 6H), 3.80 (s, 3H), 4.30 (d, 2H), 6.65 (d, 2H), 7.05 (d, 2H), 7.11 (d, 2H), 7.25 (d, 2H), 7.46 (d, 2H), 7.60 (d, 2H), 8.47 (d, 1H), 8.61 (t, 1H), 9.78 (s, 1H)	488
89	4-(3-Aminopropyl) morpholine	1.58-1.71 (m, 2H), 2.22-2.40 (m, 6H), 3.18-3.32 (m, 2H), 3.51-3.60 (m, 4H), 3.80 (s, 3H), 7.04 (d, 2H), 7.23 (d, 2H), 7.46 (d, 2H), 7.54 (d, 2H), 8.14-8.24 (m, 1H), 8.47 (d, 1H), 9.76 (s, 1H)	482
90	2-(Diisopropylamino) ethylamine	0.97 (d, 12H), 2.23-2.31 (m, 2H), 2.95 (sept, 2H), 3.06-3.20 (m, 2H), 3.81 (s, 3H), 7.04 (d, 2H), 7.24 (d, 2H), 7.46 (d, 2H), 7.56 (d, 2H), 8.07 (t, 1H), 8.47 (d, 1H), 9.77 (s, 1H)	482

Examples 91-98

The following compounds were prepared by an analogous method to that described in Example 59, using the appropriate 5-bromo-2-(4-carboxyanilino)-4-(2-pyridylamino) pyrimidine (Methods 30-31) and the appropriate amine:




5

Ex	R ¹	R ² (Precursor amine)	MS (MH ⁺)	HPLC(RT)
91	Me	1-(3-Aminopropyl)-2-pyrrolidinone	524, 526	6.61
92	Me	1-(2-Aminoethyl)pyrrolidine	496, 498	5.10
93	Me	1-(2-Aminoethyl)piperidine	510, 512	7.54
94	Me	2-Isopropylaminoethylamine ¹	484, 486	3.68
95	H	1-(3-Aminopropyl)-2-pyrrolidinone	510, 512	4.00
96	H	1-(2-Aminoethyl)pyrrolidine	482, 484	3.14
97	H	1-(2-Aminoethyl)piperidine	496, 498	4.11
98	H	2-Isopropylaminoethylamine ¹	470, 472	2.80

¹ Substituent in 4-position of the 2-anilino of the resulting example is *N*-(2-isopropylaminoethyl)carbamoyl.

Examples 99-100

10 The following compounds were prepared by an analogous method to that described in Example 59, using 4-anilino-5-bromo-2-[4-(carboxymethyl)anilino]pyrimidine (Method 32) and the appropriate amine:



Ex	R (Precursor amine)	MS (MH ⁺)	HPLC (RT)
99	2-(Diisopropylamino)ethylamine	525.2, 527.2	4.81
100	4-(2-Aminoethyl)morpholine	511.0, 512.9	2.52

Example 101**4-Anilino-5-bromo-2-{4-[3-(dimethylamino)propionamido]anilino}pyrimidine**

5 A solution of dimethylamine in tetrahydrofuran (2.0M; 4 ml) was added to a solution of 4-anilino-5-bromo-2-[4-(3-chloropropionamido)anilino]pyrimidine (Method 37; 60 mg, 0.135 mmol) in DMF (0.2 ml). The solution was heated at 80°C for 1 hour, diluted with DCM (4 ml) and loaded onto a Varian Mega Bond Elut column. Elution with 0-10% 2.0M methanolic ammonia solution in DCM gave the product (5.4 mg, 9%). MS (MH⁺): 455, 457; HPLC (RT): 2.70.

Example 102**4-Anilino-5-bromo-2-{4-[3-(isopropylamino)propionamido]anilino}pyrimidine**

Using an analogous method to that described in Example 101, but starting from 4-anilino-5-bromo-2-[4-(3-chloropropionamido)anilino]pyrimidine (Method 37) and isopropylamine, the product was obtained. MS (MH⁺): 469, 471; HPLC (RT): 3.70.

Example 103**4-Anilino-2-{4-{N-[3-(imidazol-1-yl)propyl]carbamoyl}anilino}-5-methylpyrimidine**

20 4-Anilino-2-chloro-5-methylpyrimidine (Method 23; 219 mg, 1.0 mmol) was dissolved in *n*-butanol (20 ml) and methanol (4 ml). 4-{N-[3-(imidazol-1-yl)propyl]carbamoyl}aniline (Method 40; 220 mg, 0.9 mmol) and ethereal hydrogen chloride (1.0M; 2

ml, 2.0 mmol) were added and the solution was heated at 100°C for 20 hours, and then left to stand for 4 days. The solid which separated out was collected by filtration and washed with ether (20 ml) to give the product as a hydrochloride salt (128 mg, 31%). NMR: 2.0-2.1 (m, 2H), 2.2 (s, 3H), 3.2-3.3 (m, 2H), 4.25 (t, 2H), 7.35 (t, 1H), 7.45-7.6 (m, 6H), 7.7 (d, 1H), 7.75 (d, 2H), 7.85 (s, 1H), 8.0 (s, 1H), 8.65 (br t, 1H), 9.25 (s, 1H), 10.0 (br s, 1H) 11.0 (br s, 1H); MS (MH⁺): 428.2.


Example 104

4-Anilino-5-bromo-2-{4-[N-[2-(diethylamino)ethyl]carbamoyl}anilino}pyrimidine

Using an analogous method to that described in Example 103, but starting from 4-{N-[2-(diethylamino)ethyl]carbamoyl}aniline and 4-anilino-5-bromo-2-chloropyrimidine (Method 15), the product was obtained. NMR: 1.0 (t, 6H), 2.4 (m, 4H), 3.3 (m, 4H) 7.2 (t, 1H), 7.4 (t, 2H), 7.6 (t, 6H), 8.1 (bs, 1H), 8.2 (s, 1H), 8.7 (s, 1H), 9.6 (s, 1H); MS (MH⁺): 483, 485.

Examples 105-109

The following compounds were prepared by an analogous method to that described in Example 103, using 4-{N-[3-(imidazol-1-yl)propyl]carbamoyl}aniline (Method 40) and the appropriate 5-substituted 4-anilino-2-chloropyrimidine (Methods 16, 22, 24-25):




Ex	R ¹	R ²	NMR	MS (MH ⁺)
105	Br	2-CH ₂ OH	1.93 (tt, 2H), 3.20 (dt, 2H), 4.00 (t, 2H), 4.55 (s, 2H), 6.88 (d, 1H), 7.19 (d, 1H), 7.20 (dd, 1H), 7.36 (d, 1H), 7.40 (dd, 1H), 7.63 (m, 6H), 7.89 (d, 1H), 8.24 (t, 1H), 8.26 (s, 1H), 9.59 (s, 1H)	522.2, 524.2
106	Br	4-CH ₂ OH	1.94 (tt, 2H), 3.20 (dt, 2H), 4.00 (t, 2H), 4.51 (d, 2H), 5.18 (t, 1H), 6.88 (d, 1H), 7.19 (d, 1H), 7.32 (d, 2H), 7.55 (d, 2H), 7.65 (m, 5H), 8.21 (t, 1H), 8.24 (s, 1H), 8.62 (s, 1H), 9.53 (s, 1H)	522.5, 524.6
107	Br	4-F	1.94 (tt, 2H), 3.20 (dt, 2H), 4.00 (t, 2H), 6.88 (d, 1H), 7.19 (d, 1H), 7.21 (dd, 2H), 7.60 (dd, 2H), 7.63 (m, 5H), 8.24 (s, 1H), 8.26 (t, 1H), 8.73 (s, 1H), 9.55 (s, 1H)	510.4, 512.4
108	CN	H	1.94 (tt, 2H), 3.20 (dt, 2H), 4.00 (t, 2H), 6.88 (d, 1H), 7.19 (d, 1H), 7.21 (t, 1H), 7.40 (dd, 2H), 7.56 (d, 2H), 7.65 (m, 5H), 8.31 (t, 1H), 8.53 (s, 1H), 9.59 (br. s, 1H), 10.08 (s, 1H)	439
109	Cl	4-F	1.93 (tt, 2H), 3.20 (dt, 2H), 4.00 (t, 2H), 6.87 (d, 1H), 7.19 (d, 1H), 7.22 (dd, 2H), 7.65 (m, 7H), 8.17 (s, 1H), 8.26 (t, 1H), 8.96 (s, 1H), 9.57 (s, 1H)	466, 468

¹Prepared from 4-anilino-5-cyano-2-(methanesulphonyl)pyrimidine (Method 42)

Examples 110-113


- 5 The following compounds were prepared by an analogous method to that described in Example 103, using 4-{*N*-[3-(imidazol-1-yl)propyl]carbamoyl}aniline (Method 40) and the appropriate 4-anilino-5-bromo-2-chloropyrimidine (Methods 16-18, 21):



Ex	R	MS (MH ⁺)	HPLC (RT)
110	4-SMe	538, 540	3.30
111	4-Br	572, 574	4.17
112	3,4-di-F	528, 530	3.97
113	3-F	509.9, 511.7	1.79

Examples 114-117

The following compounds were prepared by an analogous method to that described in Example 103, using 4-{*N*-[3-(imidazol-1-yl)propyl]carbamoyl}aniline (Method 40) and the appropriate 5-substituted 2-chloro-4-(pyridin-2-ylamino)pyrimidine (Methods 33-36):




Ex	R ¹	R ²	NMR	MS (MH ⁺)
114	Br	H	1.9 (m, 2H), 3.2 (q, 2H), 4.0 (t, 2H), 6.9 (s, 1H), 7.2 (m, 2H), 7.65 (s, 1H), 7.75 (s, 4H), 7.8 (m, 1H), 8.2 (d, 1H), 8.3 (t, 1H), 8.4 (m, 3H), 9.8 (s, 1H)	493, 495

115	Br	Me	1.9 (m, 2H), 2.4 (s, 3H), 3.25 (q, 2H), 4.0 (t, 2H), 6.9 (s, 1H), 7.0 (d, 1H), 7.2 (s, 1H), 7.6 (s, 1H), 7.8 (m, 5H), 8.05 (d, 1H), 8.2 (s, 1H), 8.3 (t, 1H), 8.4 (s, 1H), 9.6 (s, 1H)	507, 509
116	Cl	H	1.9 (m, 2H), 3.2 (q, 2H), 4.0 (t, 2H), 6.9 (s, 1H), 7.2 (m, 2H), 7.55 (s, 1H), 7.65 (s, 4H), 7.8 (m, 1H), 8.1 (d, 1H), 8.3 (s, 2H), 8.4 (d, 1H), 8.8 (s, 1H), 9.8 (s, 1H)	449, 451
117	Cl	Me	1.9 (m, 2H), 2.4 (s, 3H), 3.2 (q, 2H), 4.0 (t, 2H), 6.9 (s, 1H), 7.1 (d, 1H), 7.2 (s, 1H), 7.6 (s, 1H), 7.8 (m, 5H), 8.0 (d, 1H), 8.3 (m, 2H), 8.5 (s, 1H), 9.8 (s, 1H)	463, 465

Examples 118-119

The following compounds were prepared by an analogous method to that described in Example 103, using 4-{*N*-[3-(imidazol-1-yl)propyl]carbamoyl}aniline (Method 40) and the appropriate 4-substituted 5-bromo-2-chloropyrimidine intermediate (Methods 44-45):



Ex	R	NMR	MS (MH ⁺)
118	4-methylthiazol-2-yl	1.9 (m, 2H), 2.2 (s, 3H), 3.2 (q, 2H), 4.0 (t, 2H), 6.7 (s, 1H), 6.9 (s, 1H), 7.2 (s, 1H), 7.6 (s, 1H), 7.8 (m, 4H), 8.3 (m, 2H), 9.4 (s, 1H)	513, 515
119	5-methylpyrazol-3-yl	1.9 (m, 2H), 2.2 (s, 3H), 3.2 (m, 2H), 4.0 (t, 2H), 6.4 (s, 1H), 6.9 (s, 1H), 7.2 (s, 1H), 7.6 (s, 1H), 7.7 (m, 4H), 8.3 (m, 2H), 8.4 (s, 1H), 12.2 (s, 1H)	496, 498

Preparation of Starting Materials

The starting materials for the Examples above are either commercially available or are readily prepared by standard methods from known materials. For example, the following reactions are an illustration, but not a limitation, of some of the starting materials used in the

5 above reactions.

Method 1**4-Anilino-2-(4-carboxyanilino)-5-chloropyrimidine**

Ethereal hydrogen chloride (1.0M; 10 ml, 10.0 mmol) was added to a solution of 4-anilino-2,5-dichloropyrimidine (Method 2; 2.41 g, 10.0 mmol) and 4-aminobenzoic acid

10 (1.10 g, 8.0 mmol) in sulpholane (10 ml) and the mixture was heated at 140°C for 3 hours. The mixture was left to cool and acetone (75 ml) was added. The insoluble solid was collected by filtration and washed with acetone (50 ml) to give the product (2.63 g, 97%). NMR: 7.25 (t, 1H), 7.4 (t, 2H), 7.55-7.65 (m, 4H), 7.7 (d, 2H), 8.35 (s, 1H), 9.6 (br s, 1H), 10.4 (br s, 1H); MS (MH⁺): 341, 343.

15

Method 2**4-Anilino-2,5-dichloropyrimidine**

A solution of 2,4,5-trichloropyrimidine (Method 3; 5.5 g, 30.0 mmol), aniline (2.79 g, 30.0 mmol) and *N,N*-diisopropylethylamine (3.87 g, 30.0 mmol) in *n*-butanol (75 ml) was

20 heated under reflux for 4 hours. Volatile material was removed by evaporation and the residue was dissolved in DCM (100 ml). The solution was washed with water (3 x 100 ml) and saturated sodium chloride solution (100 ml) and dried. Volatile material was removed by evaporation and the residue was purified by column chromatography on silica gel, eluting with 15% ethyl acetate/isohexane, to give the product as an oil which solidified on standing

25 (3.94 g, 54%). NMR: 7.2 (t, 1H), 7.4 (t, 2H), 7.6 (d, 2H), 8.4 (s, 1H), 9.45 (br s, 1H); MS (MH⁺): 240, 242, 244.

Method 3**2,4,5-Trichloropyrimidine**

30 5-Chlorouracil (10.0 g, 68.5 mmol) was dissolved in phosphorus oxychloride (60 ml) and phosphorus pentachloride (16.0 g, 77.0 mmol) was added. The mixture was heated under reflux for 16 hours, left to cool and then poured slowly into water (200 ml) with vigorous

stirring. The mixture was stirred for 1.5 hours and then ethyl acetate (250 ml) was added. The organic layer was separated and the aqueous layer was extracted with a further portion of ethyl acetate (250 ml). The combined extracts were washed with saturated sodium bicarbonate (200 ml) and saturated sodium chloride solution (200 ml), and then dried. Volatile material

- 5 was removed by evaporation and the residue was purified by column chromatography, eluting with DCM, to give the product as a yellow liquid (6.37 g, 51%). NMR (CDCl₃): 8.62 (s, 1H); MS (MH⁺): 182, 184, 186.

Method 4


10 4-Anilino-5-bromo-2-(3-carboxyanilino)pyrimidine

Using an analogous method to that described in Method 1, but starting from 4-anilino-5-bromo-2-chloropyrimidine and 3-aminobenzoic acid, the product was obtained. NMR: 7.2-7.4 (m, 4H), 7.55-7.6 (m, 3H), 7.8 (dd, 1H), 7.95 (s, 1H), 8.4 (s, 1H), 9.3 (br s, 1H), 10.2 (br s 1H); MS (MH⁺): 385, 387.

15

Methods 5-13

The following intermediates were prepared by an analogous method to that described in Method 1, using 4-aminobenzoic acid and the appropriate 5-substituted 4-anilino-2-chloropyrimidine (Methods 14-22):



20


Method	R	MS (MH ⁺)
5	4-OMe	415, 417
6	H	385, 387
7	4-F	403, 405
8	3,4-di-F	421, 423
9	4-SMe	431, 433

10	4-Br	463, 465, 467
11	3-Me	399, 401
12	3-F	403, 405
13	2-CH ₂ OH	415, 417

Methods 14-25

The following intermediates were prepared by an analogous method to that described in Method 2, using the appropriate aniline and the appropriate 5-substituted


5 2,4-dichloropyrimidine:



Method	R ¹	R ²	MS (MH ⁺) or NMR
14	Br	4-OMe	314, 316
15	Br	H	284, 286, 288
16	Br	4-F	7.22 (m, 2H), 7.55 (m, 2H), 8.42 (s, 1H), 9.32 (s, 1H)
17	Br	3,4-di-F	7.42 (m, 2H), 7.72 (m, 1H), 8.47 (s, 1H), 9.37 (s, 1H)
18	Br	4-SMe	2.50 (s, 3H), 7.27 (d, 2H), 7.47 (d, 2H) 8.42 (s, 1H), 9.23 (s, 1H)
19	Br	4-Br	360.0, 362.0, 364.0, 366.0 (MH)
20	Br	3-Me	298, 300, 302
21	Br	3-F	6.97 (m, 1H), 7.45 (m, 3H), 8.48 (s, 1H), 9.35 (s, 1H)
22	Br	2-CH ₂ OH	314, 316
23	Me	H	220.2, 222.2
24	Br	4-CH ₂ OH	314, 316, 318
25	Cl	4-F	258, 260

Methods 26-27

The following intermediates were prepared by an analogous method to that described in Method 1, using 4-aminobenzoic acid and the appropriate 5-substituted 4-phenoxy-2-chloropyrimidine (Methods 28-29):



5

Method	R	MS (MH ⁺)
26	Br	416, 418
27	F	356

Method 28**5-Bromo-2-chloro-4-(4-methoxyphenoxy)pyrimidine**

A mixture of 5-bromo-2,4-dichloropyrimidine (5.0 g, 22.0 mmol), 4-methoxyphenol (2.72 g, 22.0 mmol) and potassium carbonate (6.07 g, 44.0 mmol) in DMF (20 ml) was stirred for 24 hours. The mixture was added to water (100 ml) and the solid which separated out was collected by filtration, washed with water (50 ml) and dried under high vacuum to give the product (6.6 g, 96%). NMR: 3.8 (s, 3H), 7.0 (d, 2H), 7.2 (d, 2H), 8.8 (s, 1H).


15 **Method 29****2-Chloro-5-fluoro-4-(4-methoxyphenoxy)pyrimidine**

Using an analogous method to that described in Method 28, but starting from 2,4-dichloro-5-fluoropyrimidine and 4-methoxyphenol, the product was obtained. NMR: 3.84 (s, 3H), 6.79-7.00 (m, 2H), 7.06-7.18 (m, 2H), 8.31-8.36 (m, 1H); MS (M⁺): 254, 256.

20

Methods 30-31

The following intermediates were prepared by an analogous method to that described in Method 1, using 4-aminobenzoic acid and the appropriate 5-bromo-2-chloro-4-(2-pyridylamino)pyrimidine (Methods 32-33):



5


Method	R	MS (MH ⁺)
30	Me	400, 402
31	H	386, 388

Method 32**4-Anilino-5-bromo-2-[4-(carboxymethyl)anilino]pyrimidine**

A mixture of 4-anilino-5-bromo-2-chloropyrimidine (Method 15; 1.50 g, 5.3 mmol), 4-aminophenylacetic acid (0.76 g, 5.0 mmol) and concentrated hydrochloric acid (0.98 ml, 5.3 mmol) in n-butanol (20 ml) was heated at 100°C for 18 hours. The solid which separated out on cooling was collected by filtration and washed with n-butanol (20 ml) and diethyl ether (20 ml). The solid was dissolved in THF (20 ml) and methanol (10 ml). 2M Sodium hydroxide (3.80 ml, 7.6 mmol) was added and the solution was stirred for 18 hours. Volatile material was removed by evaporation and the residue was partitioned between ether (20 ml) and water (20 ml). The aqueous phase was separated and acidified to pH 3 to give the product as a white solid (350 mg). NMR: 7.0 (d, 2H), 7.2 (m, 1H), 7.4 (m, 2H), 7.5 (d, 2H), 7.6 (d, 2H), 8.2 (s, 1H), 8.6 (s, 1H), 9.3 (s, 1H); MS (MH⁺); 399, 401.

20 Methods 33-36

The following intermediates were prepared by an analogous method to that described in Method 2, using the appropriate 2-aminopyridine and the appropriate 5-substituted 2,4-dichloropyrimidine:



Method	R¹	R²	MS (MH⁺)
33	Br	Me	299, 301
34	Br	H	285, 287
35	Cl	H	240, 242
36	Cl	Me	254, 256

Method 37**4-Anilino-5-bromo-2-[4-(3-chloropropionamido)anilino]pyrimidine**

5 Chloropropionyl chloride (0.143 g, 1.13 mmol) was added to a stirred solution of 2-(4-aminoanilino)-4-anilino-5-bromopyrimidine (Method 38; 0.40 g, 1.13 mmol) and triethylamine (0.173 ml, 1.24 mmol) in DMF (1 ml). The mixture was stirred at room temperature and then water (4 ml) was added. The precipitated solid was collected by filtration and washed with ether (1 ml) to give the product (0.21 g, 42%) as a brown solid. MS
10 (MH⁺): 448, 450.

Method 38**2-(4-Aminoanilino)-4-anilino-5-bromopyrimidine**

Sodium hydrosulphite (9.5 g, 54.5 mmol) was added over 30 minutes to a stirred
15 solution of anilino-5-bromo-2-(4-nitroanilino)pyrimidine (Method 39; 7.0 g, 18.2 mmol) in a hot mixture of ethanol and water (1:1, 100 ml). The resulting suspension was stirred for a further two hours and insoluble material was removed by filtration. The filtrate was concentrated and the residue was partitioned between 1M aqueous sodium hydroxide (50 ml), and DCM (100 ml). The organic layer was separated, washed with water (2 x 10 ml) and
20 dried. Volatile material was removed by evaporation to give the product (4.55 g, 70%). MS (MH⁺): 356, 358.

Method 39**4-Anilino-5-bromo-2-(4-nitroanilino)pyrimidine**

Using an analogous method to that described in Method 1, but starting from 4-anilino-5-bromo-2-chloropyrimidine (Method 15), the product was obtained. MS (MH⁺):
5 386, 388.

Method 40**4-{N-[3-(Imidazol-1-yl)propyl]carbamoyl}aniline**

A mixture of 4-{N-[3-(imidazol-1-yl)propyl]carbamoyl}nitrobenzene (Method 41; 2.0
10 g, 7.30 mmol), 10% palladium on carbon (100 mg) cyclohexene (40 ml) and ethanol (80 ml) was heated under reflux under an atmosphere of nitrogen for 4 hours. Two further portions of cyclohexene (40 ml) were added and heating was continued for 1 hour after each addition. The mixture was left to cool and the catalyst was removed by filtration. The filtrate was concentrated by evaporation and the residue was recrystallized from a mixture of ethanol and
15 ether to give the product (1.2 g, 67%). NMR: 1.9-2.0 (m, 2H), 3.1-3.2 (m, 2H), 4.0 (t, 2H), 5.6 (br s 2H), 6.55 (d, 2H), 6.85 (s, 1H), 7.2 (s, 1H), 7.55 (d, 2H), 7.65 (s, 1H), 8.0 (br t, H).

Method 41**4-{N-[3-(Imidazol-1-yl)propyl]carbamoyl}nitrobenzene**

20 A solution of 4-nitrobenzoyl chloride (18.5 g, 0.1 mol) in DCM (200 ml) was added dropwise to a stirred solution of 1-(3-aminopropyl)imidazole (14.0 g, 0.112 mol) and triethylamine (15 ml, 0.107 mol) in DCM (200 ml) at 0°C. The mixture was stirred at 0°C for 1 hour, and the solid which had separated out was collected by filtration and washed with water (100 ml) and acetone (100 ml) to give the product (21.6 g, 79%). NMR: 1.9-2.0 (m,
25 2H), 3.2-3.3 (m, 2H), 4.0 (t, 2H), 6.85 (s, 1H), 7.2 (s, 1H), 7.65 (s, 1H), 8.05 (d, 2H), 8.3 (d, 2H), 8.8 (br s, 1H).

Method 42**4-Anilino-5-cyano-2-(methanesulphonyl)pyrimidine**

30 3-Chloroperoxybenzoic acid (57-86%; 2.67 g, 8.8-13.3 mmol) was added in aliquots to a solution of 4-anilino-5-cyano-2-(methylthio)pyrimidine (Method 43; 1.0 g, 4.13 mmol) in chloroform (100 ml), and the mixture was stirred for 2 hours. The mixture was washed with

saturated sodium bicarbonate (100 ml), water (100 ml), and saturated sodium chloride (100 ml) and dried. Volatile material was removed by evaporation and the residue was taken up in DCM (10 ml). The solution was loaded onto a silica column pre-equilibrated with 20% ethyl acetate solution in isohexane. Elution with 20-50% ethyl acetate in isohexane and

- 5 concentration of the appropriate fractions gave the product as a yellow solid (680 mg, 61%).
NMR (CDCl₃): 3.26 (s, 3H), 7.30 (t, 1H), 7.44 (dd, 2H), 7.57 (d, 2H), 7.65 (br s, 1H), 8.71 (s, 1H); MS (MH⁺): 274.9.

Method 43


10 4-Anilino-5-cyano-2-(methylthio)pyrimidine

Using a method analogous to that described in Method 2, but starting from 4-chloro-5-cyano-2-(methylthio)pyrimidine (obtained as described in J. Het. Chem. 1971, 8, 445) and performing the reaction at 85°C, the product was obtained. NMR (CDCl₃): 2.51 (s, 3H), 7.15 (br s, 1H), 7.20 (t, 1H), 7.40 (dd, 2H), 7.57 (d, 2H), 8.38 (s, 1H).

15

Methods 44-45

The following intermediates were prepared by an analogous method to that described in Method 2, using the appropriate amino heterocycle and 5-bromo-2,4-dichloropyrimidine:



Method	R	MS (MH ⁺)
44	4-methylthiazol-2-yl	303, 305
45	5-methylpyrazol-3-yl	288, 290

20

Example 120

The following illustrate representative pharmaceutical dosage forms containing the compound of formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof (hereafter compound X), for therapeutic or prophylactic use in humans:-

(a): Tablet I	mg/tablet
Compound X	100
Lactose Ph.Eur	182.75
Croscarmellose sodium	12.0
Maize starch paste (5% w/v paste)	2.25
Magnesium stearate	3.0

(b): Tablet II	mg/tablet
Compound X	50
Lactose Ph.Eur	223.75
Croscarmellose sodium	6.0
Maize starch	15.0
Polyvinylpyrrolidone (5% w/v paste)	2.25
Magnesium stearate	3.0

(c): Tablet III	mg/tablet
Compound X	1.0
Lactose Ph.Eur	93.25
Croscarmellose sodium	4.0
Maize starch paste (5% w/v paste)	0.75
Magnesium stearate	1.0

(d): Capsule	mg/capsule
Compound X	10
Lactose Ph.Eur	488.5
Magnesium stearate	1.5

(e): Injection I	(50 mg/ml)
Compound X	5.0% w/v
1M Sodium hydroxide solution	15.0% v/v
0.1M Hydrochloric acid	(to adjust pH to 7.6)
Polyethylene glycol 400	4.5% w/v
Water for injection	to 100%

(f): Injection II	10 mg/ml
Compound X	1.0% w/v
Sodium phosphate BP	3.6% w/v
0.1M Sodium hydroxide solution	15.0% v/v
Water for injection	to 100%


(g): Injection III	(1mg/ml, buffered to pH6)
Compound X	0.1% w/v
Sodium phosphate BP	2.26% w/v
Citric acid	0.38% w/v
Polyethylene glycol 400	3.5% w/v
Water for injection	to 100%

5 Note

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.

Claims


1. A pyrimidine derivative of the formula (I):




(I)

wherein:

Q_1 and Q_2 are independently selected from aryl or carbon linked heteroaryl; and one of Q_1 and Q_2 or both of Q_1 and Q_2 is substituted on a ring carbon by one substituent of the formula (Ia) or (Ia'):



(Ia)



(Ia')

wherein:

Y is -NHC(O)- or -C(O)NH-;

Z is R^aO- , R^bR^cN- , R^dS- , $R^eR^fNNR^g-$, C_{3-8} cycloalkyl, phenyl or a heterocyclic group;

- 15 wherein said phenyl, C_{3-8} cycloalkyl or heterocyclic group are optionally substituted on a ring carbon by one or more groups selected from R^h ; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^i ;

- 20 R^a , R^b , R^c , R^d , R^e , R^f and R^g are independently selected from hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl and C_{3-8} cycloalkyl; wherein said C_{1-4} alkyl, C_{2-4} alkenyl and C_{3-8} cycloalkyl are optionally substituted by one or more groups selected from R^j ;

n is 0 or 1;

m is 1, 2 or 3;

- 25 Q_3 is a nitrogen linked heterocycle; wherein said heterocycle is optionally substituted on a ring carbon by one or more groups selected from R^k ; and wherein if said heterocyclic

group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^m;

G is -O- or -NR²-;

R² is selected from hydrogen, C₁₋₆alkyl, C₃₋₆alkenyl and C₃₋₆alkynyl; wherein said

- 5 C₁₋₆alkyl, C₃₋₆alkenyl and C₃₋₆alkynyl are optionally substituted by one or more groups selected from Rⁿ;

- R¹ is selected from hydrogen, halo, hydroxy, nitro, amino, *N*-(C₁₋₃alkyl)amino, *N,N*-di-(C₁₋₃alkyl)amino, cyano, trifluoromethyl, trichloromethyl, C₁₋₃alkyl [optionally substituted by 1 or 2 substituents independently selected from halo, cyano, amino, *N*-(C₁₋₃alkyl)amino, *N,N*-di-(C₁₋₃alkyl)amino, hydroxy and trifluoromethyl], C₃₋₅alkenyl [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent], C₃₋₅alkynyl, C₁₋₃alkoxy, mercapto, C₁₋₃alkylsulphanyl, carboxy and C₁₋₃alkoxycarbonyl;

- Q₁ is optionally substituted on a ring carbon by one to four substituents independently selected from halo, mercapto, nitro, formyl, formamido, carboxy, cyano, amino, ureido, carbamoyl, sulphamoyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl [wherein said C₁₋₄alkyl, C₂₋₄alkenyl and C₂₋₄alkynyl are optionally substituted by one or more groups selected from R^o], C₁₋₄alkanoyl, C₁₋₄alkoxycarbonyl, heterocyclic group, C₁₋₄alkylS(O)_a wherein a is 0 to 2 [optionally substituted by hydroxy], *N'*-(C₁₋₄alkyl)ureido, *N',N'*-di-(C₁₋₄alkyl)ureido, *N'*-(C₁₋₄alkyl)-*N*-(C₁₋₄alkyl)ureido, *N',N'*-di-(C₁₋₄alkyl)-*N*-(C₁₋₄alkyl)ureido, *N*-C₁₋₄alkylamino, *N,N*-di-(C₁₋₄alkyl)amino, *N*-(C₁₋₄alkyl)sulphamoyl, *N,N*-di-(C₁₋₄alkyl)sulphamoyl, *N*-C₁₋₄alkylcarbamoyl, *N,N*-di-(C₁₋₄alkyl)carbamoyl and C₁₋₄alkanoylamino;

- and also independently, or in addition to, the above substituents, Q₁ may be optionally substituted by one to two substituents independently selected from aryl, C₃₋₈cycloalkyl and a heterocyclic group; wherein said aryl, C₃₋₈cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^p; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^q;

and also independently, or in addition to, the above substituents, Q₁ may be optionally substituted by one C₁₋₄alkoxy or by one hydroxy substituent;

- 30 Q₂ is optionally substituted on a ring carbon by one to four substituents independently selected from halo, hydroxy, mercapto, nitro, formyl, formamido, carboxy, cyano, amino, ureido, carbamoyl, sulphamoyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy [wherein said

C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl and C₁₋₄alkoxy are optionally substituted by one or more groups selected from R^j, C₁₋₄alkanoyl, C₁₋₄alkoxycarbonyl, heterocyclic group, C₁₋₄alkylS(O)_a wherein a is 0 to 2 [optionally substituted by hydroxy], N^j-(C₁₋₄alkyl)ureido, N^j,N^j-di-(C₁₋₄alkyl)ureido, N^j-(C₁₋₄alkyl)-N-(C₁₋₄alkyl)ureido, N^j,N^j-di-(C₁₋₄alkyl)-N-(C₁₋₄alkyl)ureido, N-C₁₋₄alkylamino, N,N-di-(C₁₋₄alkyl)amino, N-(C₁₋₄alkyl)sulphamoyl, N,N-di-(C₁₋₄alkyl)sulphamoyl, N-C₁₋₄alkylcarbamoyl, N,N-di-(C₁₋₄alkyl)carbamoyl, C₂₋₄alkenyloxy, C₂₋₄alkynyloxy, C₁₋₄alkanoylamino and a group of formula (Ia) or (Ia') as depicted above;

and also independently, or in addition to, the above substituents, Q₂ may be optionally substituted by one to two substituents independently selected from aryl, C₃₋₈cycloalkyl or a heterocyclic group; wherein said aryl, C₃₋₈cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^s; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^t;

R^j, R^a, R^o and R^r are independently selected from hydroxy, halo, amino, cyano, formyl, formamido, carboxy, nitro, mercapto, carbamoyl, sulphamoyl, N-C₁₋₄alkylamino, N,N-di-(C₁₋₄alkyl)amino, C₁₋₄alkanoyl, C₁₋₄alkanoyloxy, C₁₋₄alkoxy, C₁₋₄alkoxycarbonyl, N-C₁₋₄alkylcarbamoyl, N,N-di-(C₁₋₄alkyl)carbamoyl, C₁₋₄alkanoylamino, C₁₋₄alkylS(O)_a wherein a is 0 to 2, C₁₋₄alkylsulphonylamino, N-(C₁₋₄alkyl)sulphamoyl, N-(C₁₋₄alkyl)₂sulphamoyl, N-(C₁₋₄alkyl)carbamoyl, N-(C₁₋₄alkyl)₂carbamoyl, phenyl, phenylthio, phenoxy, C₃₋₈cycloalkyl and a heterocyclic group; wherein said phenyl, phenylthio, phenoxy, C₃₋₈cycloalkyl or heterocyclic group may be optionally substituted on a ring carbon by one or more groups selected from R^u; and wherein if said heterocyclic group contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from R^v;

R^b, R^k, R^p, R^s and R^u are independently selected from hydroxy, halo, amino, cyano, formyl, formamido, carboxy, nitro, mercapto, carbamoyl, sulphamoyl, C₁₋₄alkyl [optionally substituted by one or more groups selected from halo, cyano, amino, N-C₁₋₄alkylamino, N,N-di-(C₁₋₄alkyl)amino or hydroxy], C₂₋₄alkenyl [optionally substituted by one or more groups selected from halo], C₂₋₄alkynyl, N-C₁₋₄alkylamino, N,N-di-(C₁₋₄alkyl)amino, C₁₋₄alkanoyl, C₁₋₄alkanoyloxy, C₁₋₄alkoxy [optionally substituted by one or more groups selected from halo], C₁₋₄alkoxycarbonyl, N-C₁₋₄alkylcarbamoyl, N,N-di-(C₁₋₄alkyl)carbamoyl,

C₁₋₄alkanoylamino, C₁₋₄alkylS(O)_a wherein a is 0 to 2, C₁₋₄alkylsulphonylamino, N-(C₁₋₄alkyl)sulphamoyl, N-(C₁₋₄alkyl)₂sulphamoyl, phenyl, C₃₋₈cycloalkyl and a heterocyclic group; and

Rⁱ, R^q, Rⁱ and R^v are independently selected from C₁₋₄alkyl, C₁₋₄alkanoyl,

- 5 C₁₋₄alkylsulphonyl, C₁₋₄alkoxycarbonyl, carbamoyl, N-(C₁₋₄alkyl)carbamoyl, N,N-(C₁₋₄alkyl)carbamoyl, benzyl, benzyloxycarbonyl, benzoyl and phenylsulphonyl; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

2. A pyrimidine derivative according to claim 1 wherein Q₁ is phenyl or a
10 pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

3. A pyrimidine derivative according either of claims 1 or 2 wherein Q₂ is phenyl, pyridyl, thiazolyl or pyrazolyl or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

15

4. A pyrimidine derivative according any one of claims 1 to 3 wherein the substituent (Ia) or (Ia') is N-[2-(3-aza-8-oxabicyclo[3,2,1]hex-3-yl)ethyl]carbamoyl, N-(2-di-*n*-butylaminoethyl)carbamoyl, N-(2-diethylaminoethyl)carbamoyl, N-(2-diisopropylaminoethyl)carbamoyl, N-[2-(3,5-dimethylpyrazol-1-yl)ethyl]carbamoyl,
20 N-(2-indol-3-ylethyl)carbamoyl, N-(2-isopropylaminoethyl)carbamoyl, N-[2-(2-methyl-5-nitroimidazol-1-yl)ethyl]carbamoyl, N-(2-methylthioethyl)carbamoyl, N-(2-morpholinoethyl)carbamoyl, N-(2-piperidin-1-ylethyl)carbamoyl, N-(2-pyrrolidin-1-ylethyl)carbamoyl, N-(2-thien-2-ylethyl)carbamoyl, N-(2-thiomorpholinoethyl)carbamoyl, N-(3-*n*-butoxypropyl)carbamoyl,
25 N-(3-di-*n*-butylaminopropyl)carbamoyl, N-(3-imidazol-1-ylpropyl)carbamoyl, N-[3-(4-methylpiperazin-1-yl)propyl]carbamoyl, N-(3-methylthiopropyl)carbamoyl, N-(3-morpholinopropyl)carbamoyl, N-[3-(2-oxypyrrolidin-1-yl)propyl]carbamoyl, N-(3-pyrrolidin-1-ylpropyl)carbamoyl, N-(2-di-isopropylaminoethyl)carbamoylmethyl, N-(2-morpholinoethyl)carbamoylmethyl, N-(4-dimethylaminobenzylamine)carbamoyl,
30 N-(imidazo[1,2-*a*]pyrid-2-ylmethyl)carbamoyl, N-(5-methylfur-2-ylmethyl)carbamoyl, 4-(pyrrolidin-1-yl)piperidin-1-ylcarbonyl, 3-(dimethylamino)propanamide or

3-(isopropylamino)propanamide or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

5. A pyrimidine derivative according any one of claims 1 to 4 wherein the substituent of
5 formula (Ia) or (Ia') is on ring Q₁ or a pharmaceutically acceptable salt or *in vivo*
hydrolysable ester thereof.

6. A pyrimidine derivative according any one of claims 1 to 5 wherein when Q₁ is phenyl
the substituent of formula (Ia) or (Ia') is in either the para- or meta- position relative to the
10 -NH- or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

7. A pyrimidine derivative according any one of claims 1 to 6 wherein G is -O- or -NH-
or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

15 8. A pyrimidine derivative according any one of claims 1 to 7 wherein R¹ is fluoro,
chloro, bromo, methyl or cyano or a pharmaceutically acceptable salt or *in vivo* hydrolysable
ester thereof.


9. A pyrimidine derivative according any one of claims 1 to 8 wherein Q₁ is
20 unsubstituted except by the substituent of formula (Ia) or (Ia') or a pharmaceutically
acceptable salt or *in vivo* hydrolysable ester thereof.

10. A pyrimidine derivative according any one of claims 1 to 9 wherein Q₂ is
unsubstituted or substituted by one or two groups selected from fluoro, bromo, methyl,
25 methoxy, methylthio or hydroxymethyl or a pharmaceutically acceptable salt or *in vivo*
hydrolysable ester thereof.

11. A pyrimidine derivative according any one of claims 1 to 10 selected from:
2-{4-[N-(3-imidazol-1-ylpropyl)carbamoyl]anilino}-4-anilino-5-bromopyrimidine;
30 2-{4-[N-(2-pyrrolidin-1-ylethyl)carbamoyl]anilino}-4-(3-fluoroanilino)-5-bromopyrimidine;
2-{4-[N-(2-isopropylaminoethyl)carbamoyl]anilino}-4-(3-fluoroanilino)-5-bromopyrimidine;


- 2-{4-[N-(2-pyrrolidin-1-ylethyl)carbamoyl]anilino}-4-(2-hydroxymethylanilino)-5-bromopyrimidine;
- 2-{4-[N-(2-isopropylaminoethyl)carbamoyl]anilino}-4-(2-hydroxymethylanilino)-5-bromopyrimidine;
- 5 2-{4-[N-(2-pyrrolidin-1-ylethyl)carbamoyl]anilino}-4-(6-methylpyrid-2-ylamino)-5-bromopyrimidine;
- 2-{4-[N-(2-isopropylaminoethyl)carbamoyl]anilino}-4-(6-methylpyrid-2-ylamino)-5-bromopyrimidine;
- 2-{4-[N-(2-pyrrolidin-1-ylethyl)carbamoyl]anilino}-4-(pyrid-2-ylamino)-5-bromopyrimidine;
- 10 2-{4-[N-(3-imidazol-1-ylpropyl)carbamoyl]anilino}-4-(pyrid-2-ylamino)-5-bromopyrimidine;
- 2-{4-[N-(3-imidazol-1-ylpropyl)carbamoyl]anilino}-4-(6-methylpyrid-2-ylamino)-5-bromopyrimidine;
- or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

- 15 12. A process for preparing a compound of formula (I) or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof as claimed in any one of claims 1 to 11 which is:
- a) for compounds of formula (I) where G is -NR²-; reacting a pyrimidine of formula (II):



(II)


- 20 wherein L is a displaceable group as defined below, with a compound of formula (III):



(III)


where G is -NR²-;

- b) reaction of a pyrimidine of formula (IV):



(IV)


wherein L is a displaceable group as defined below, with a compound of formula (V):



(V)


5

c) for compounds of formula (I) wherein the sidechain is of formula (Ia) and Y is -C(O)NH-; by reaction of an acid of formula (VI):




(VI)

10 or an activated derivative thereof; with an amine of formula (VII):



(VII)


d) for compounds of formula (I) wherein the sidechain is of formula (Ia) and Y is -NHC(O)- by reaction of an amine of formula (VIII):



(VIII)

15


with an acid of formula (IX):



(IX)

20 or an activated derivative thereof;

e) for compounds of formula (I) wherein the sidechain is of formula (Ia'); by reaction of an acid of formula (VI) (or an activated derivative thereof) with an amine of formula (X):



5 and thereafter if necessary:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

10 13. A pharmaceutical composition which comprises a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1 to 11, in association with a pharmaceutically acceptable diluent or carrier.

14. A pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1 to 11, for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

15. A pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1 to 11, for use as a
20 medicament.

16. The use of a pyrimidine derivative of the formula (I), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1 to 11, in the manufacture of a medicament for use in the production of an anti-cancer, cell cycle inhibitory
25 (anti-cell-proliferation) effect and/or a FAK inhibitory (anti-cell migration and/or apoptosis inducing) effect in a warm-blooded animal such as man.

17. A method for producing an anti-cancer, cell cycle inhibitory (anti-cell-proliferation) effect and/or a FAK inhibitory (anti-cell migration and/or apoptosis inducing) effect in a
30 warm-blooded animal, such as man, in need of such treatment which comprises administering

to said animal an effective amount of a pyrimidine derivative as claimed in any one of claims 1 to 11, or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 01/00824

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D239/48 C07D403/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 09852 A (CIBA GEIGY AG) 13 April 1995 (1995-04-13) claims	4-17
A	EP 0 564 409 A (CIBA GEIGY AG) 6 October 1993 (1993-10-06) claims	4-17
P, Y	WO 00 53595 A (BREAUULT GLORIA ANNE ; JAMES STEWART RUSSELL (GB); PEASE JANE ELIZAB) 14 September 2000 (2000-09-14) claims; examples	4-17
P, Y	WO 00 12485 A (BREAUULT GLORIA ANNE ; PEASE JANET ELIZABETH (GB); ZENECA LTD (GB)) 9 March 2000 (2000-03-09) claims; examples	4-17
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

1 June 2001

Date of mailing of the international search report

12/06/2001

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Menegaki, F

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 01/00824

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	WO 00 39101 A (BREault GLORIA ANNE ;PEASE JANET ELIZABETH (GB); ASTRAZENECA UK LT) 6 July 2000 (2000-07-06) claims -----	4-17

INTERNATIONAL SEARCH REPORT

International application No.

PCT/ GB/ 01/ 00824

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

SEE ANNEXE

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-3

Present claims 1-3 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds according to claims 4-16 and the specific examples in the description, wherein groups Q1/Q2/Q3 and substituents thereof are clearly defined.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/00824

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9509852 A	13-04-1995	AU 693804 B	09-07-1998
		AU 7697594 A	01-05-1995
		CA 2148477 A	13-04-1995
		EP 0672040 A	20-09-1995
		JP 8504834 T	28-05-1996
		US 5521184 A	28-05-1996
		US 5543520 A	06-08-1996
EP 0564409 A	06-10-1993	AT 188964 T	15-02-2000
		AU 3569493 A	07-10-1993
		BR 1100739 A	06-06-2000
		CA 2093203 A	04-10-1993
		CN 1077713 A,B	27-10-1993
		CZ 9300560 A	16-02-1994
		DE 59309931 D	24-02-2000
		DK 564409 T	19-06-2000
		ES 2142857 T	01-05-2000
		FI 931458 A	04-10-1993
		GR 3032927 T	31-07-2000
		HU 64050 A	29-11-1993
		IL 105264 A	11-04-1999
		JP 2706682 B	28-01-1998
		JP 6087834 A	29-03-1994
		MX 9301929 A	29-07-1994
		NO 302473 B	09-03-1998
		NZ 247299 A	26-07-1995
		PT 564409 T	30-06-2000
		RU 2125992 C	10-02-1999
		SG 43859 A	14-11-1997
		SK 28093 A	06-04-1994
		US 5521184 A	28-05-1996
		ZA 9302397 A	04-10-1993
WO 0053595 A	14-09-2000	AU 2818700 A	28-09-2000
WO 0012485 A	09-03-2000	AU 5438299 A	21-03-2000
WO 0039101 A	06-07-2000	AU 1874300 A	31-07-2000